arXiv:1409.3872v1 [math.DG] 12 Sep 2014

Minimal two-spheres of low index in manifolds with positive complex sectional curvature

John Douglas Moore and Robert Ream
Department of Mathematics
University of California
Santa Barbara, CA, USA 93106

September 16, 2014

Abstract

Suppose that S^n is given a Riemannian metric of positive complex sectional curvatures for which the minimal two-spheres of Morse index $\leq 2n-5$ are all Morse nondegenerate. Then the number of minimal two spheres of Morse index k, for $n-2 \leq k \leq 2n-5$, is at least $p_3(k-n+2)$, where $p_3(k-n+2)$ is the number of $(k-n+2)$-cells in the Schubert cell decomposition for $G_3(\mathbb{R}^{n+1})$.

1 Introduction

Suppose that M is a compact Riemannian manifold, $\mathrm{Map}(S^2, M)$ the space of smooth maps from the Riemann sphere S^2 to M. An element $f \in \mathrm{Map}(S^2, M)$ is *harmonic* if it is critical point for the energy function

$$E : \mathrm{Map}(S^2, M) \longrightarrow \mathbb{R},$$

which is defined by the Dirichlet integral

$$E(f) = \frac{1}{2}\int_{S^2} |df|^2 dA = \frac{1}{2}\int_{S^2} \left[\left| \frac{\partial f}{\partial x} \right|^2 + \left| \frac{\partial f}{\partial y} \right|^2 \right] dxdy.$$

In the middle integrand of this formula, $|df|$ and dA are calculated with respect to the standard Riemannian metric of constant curvature and total area one on S^2, while in the right-hand integrand (x, y) are isothermal coordinates on S^2 with respect to the standard conformal structure on the Riemann sphere S^2, the integrand being independent of the choice. If the Riemannian metric on M is induced by an imbedding of $M \subseteq \mathbb{R}^N$ into Euclidean space (always possible by the Nash imbedding theorem), the Euler-Lagrange equation in isothermal parameters for the energy function is

$$\left(\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} \right)^{\top} = 0,$$

where $(\cdot)^\top$ represents orthogonal projection from $\mathbb{R}^N$ into the tangent space. It is well-known that harmonic maps from the Riemann sphere are also critical points for the area function, and hence also represent parametrized *minimal* two-spheres, possibly with branch points.

The energy function E is invariant under the group of linear fractional transformations: if we regard S^2 as the one-point compactification $\mathbb{C} \cup \{\infty\}$ of the complex plane $\mathbb{C}$ with the standard coordinate $z = x + iy$, and if $T : S^2 \to S^2$ is the linear fractional transformation (a conformal diffeomorphism) defined by

$$T(z) = \frac{az+b}{cz+d}, \quad \text{for} \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2,\mathbb{C}),$$

then $E(f \circ T) = E(f)$. Note that $-I \in SL(2,\mathbb{C})$ acts as the identity, so the group of linear fractional transformations is actually

$$\mathcal{G} = PSL(2,\mathbb{C}) = \frac{SL(2,\mathbb{C})}{\pm I},$$

and if we want to allow orientation-reversing automorphisms of S^2 we must replace $PSL(2,\mathbb{C})$ by the larger group

$$\widehat{\mathcal{G}} = PSL(2,\mathbb{C}) \cup R \cdot PSL(2,\mathbb{C}),$$

where R is some anticonformal reflection through a great circle. A nonconstant minimal two-sphere is *prime* if it is not a branched cover of a minimal two-sphere of smaller area, and two prime minimal surfaces are *geometrically distinct* if they do not lie on the same $\widehat{\mathcal{G}}$-orbit.

Our goal here is to develop the Morse theory techniques needed to estimate the number of geometrically distinct prime minimal two-spheres of low Morse index, using a perturbation of the energy proposed by Sacks and Uhlenbeck [32]. This perturbation is the α-*energy*, for $\alpha > 1$, and it is defined by

$$E_\alpha : \mathrm{Map}(S^2, M) \longrightarrow \mathbb{R}, \quad E_\alpha(f) = \frac{1}{2}\int_{S^2} (1 + |df|^2)^\alpha dA - \frac{1}{2}, \tag{1}$$

where $|df|$ and dA are again defined in terms of the canonical Riemannian metric of constant curvature and total area one on S^2, there being exactly one such metric up to diffeomorphisms isotopic to the identity. As explained by Sacks and Uhlenbeck, the α-energy satisfies Condition C of Palais and Smale on the completion $L_1^{2\alpha}(S^2, M)$ of $\mathrm{Map}(S^2, M)$ with respect to the $L_1^{2\alpha}$ Sobolev norm, and $L_1^{2\alpha}(S^2, M) \subseteq C^0(S^2, M)$ when $\alpha > 1$. These facts make it possible to develop a Liusternik-Schnirelman theory for E_α and a Morse theory for further perturbations of E_α which have Morse nondegenerate critical points. Note that $E_\alpha(f) \to E(f)$ as $\alpha \to 1$, and we want to take the limit as $\alpha \to 1$.

There are two problems with developing a partial Morse theory for the energy E itself:

Bubbling. As observed by Sacks and Uhlenbeck, a subsequence of α-energy critical points will not necessarily converge to a single minimal surface as $\alpha \to 1$,

but instead may converge to a bubble tree, consisting of several minimal two-spheres connected by necks. (Since some of the limiting harmonic two-spheres must be nonconstant, Sacks and Uhlenbeck were able to prove the existence of at least one nonconstant minimal two-sphere in any compact simply connected Riemannian manifold but their method by itself does not determine the Morse index of the minimal two-spheres obtained.)

Branched covers. Branched covers of degree d of a given prime minimal two-sphere form a critical submanifold of real dimension $4d+2$, as we will see. But in general the space of branched covers is not a nondegenerate critical submanifold in the sense of Bott, so it is not straightforward to calculate its contribution to the topology of $\mathrm{Map}(S^2, M)$.

When these two difficulties can be controlled, one can expect theorems similar to those for closed geodesics in Riemannian manifolds.

For example, a theorem of Ballmann, Thorbergsson and Ziller on closed geodesics [2] states that a generic simply connected compact Riemannian manifold whose real sectional curvatures satisfy the inequality $1/4 < K(\sigma) \leq 1$ must contain at least $p_2(\lambda - n + 1)$ geometrically distinct smooth closed geodesics of Morse index λ, where $p_2(k)$ is the number of k-cells in the Schubert cell decomposition of $G_2(\mathbb{R}^{n+1})$. Their result can be obtained by establishing equivariant Morse inequalities for the action function on the space $\mathrm{Map}(S^1, M)$ of loops in M. (The reader can refer to [26] for a discussion of the Schubert decomposition of $G_m(\mathbb{R}^N)$ used in the argument.) We will prove an analog of this theorem (with a stronger pinching constant) for minimal two-spheres when the Morse index is sufficiently small:

Theorem 1.1. *Suppose that the compact smooth simply connected manifold M has dimension $n \geq 4$, and has a Riemannian metric whose real sectional curvatures $K_r(\sigma)$ satisfy the pinching condition*

$$\frac{1}{2} < K_r(\sigma) \leq 1. \tag{2}$$

If all prime minimal two-spheres of Morse index $\leq 2n-5$ in M are immersions and lie on Morse nondegenerate critical submanifolds, each a $\widehat{\mathcal{G}}$-orbit, where $\widehat{\mathcal{G}}$ is the group of conformal and anticonformal automorphisms of S^2, then the number of geometrically distinct prime minimal two spheres of Morse index λ is at least $p_3(\lambda - n + 2)$ for $n - 2 \leq \lambda \leq 2n - 5$, where $p_3(k)$ is the number of k-cells in the Schubert cell decomposition for $G_3(\mathbb{R}^{n+1})$.

In fact, we will weaken the pinching condition (2) somewhat by making use of complex sectional curvatures. We recall that in normal coordinates $(x^1, \ldots, x^n)$ centered at a point p on M, the Riemannian metric can be expressed by a Taylor series

$$g_{ij} = \delta_{ij} - \frac{1}{3} \sum_{k,l} R_{ikjl}(p) x^k x^l + \text{ higher order terms},$$

and the curvature operator can be defined to be the linear map $\mathcal{R} : \Lambda^2 T_p M \longrightarrow$

$\Lambda^2 T_pM$ such that

$$\mathcal{R}\left(\left.\frac{\partial}{\partial x^i}\right|_p \wedge \left.\frac{\partial}{\partial x^j}\right|_p\right) = \sum_{k,l} R_{ijkl}(p)\left.\frac{\partial}{\partial x^k}\right|_p \wedge \left.\frac{\partial}{\partial x^l}\right|_p.$$

If z and w are linearly independent elements of $T_pM \otimes \mathbb{C}$, the *complex sectional curvature* of the two-plane σ spanned by z and w is

$$K(\sigma) = \frac{\langle \mathcal{R}(z \wedge w), \bar{z} \wedge \bar{w}\rangle}{\langle z \wedge w, \bar{z} \wedge \bar{w}\rangle}, \tag{3}$$

where the curvature operator $\mathcal{R}$ and the inner product have been extended complex linearly and the bar denotes complex conjugation. The complex sectional curvature arises quite naturally in the formula for second variation of energy for a minimal two-sphere f, as presented in [22]. This second variation is the symmetric bilinear form

$$d^2E(f) : T_f\mathrm{Map}(S^2, M) \otimes \mathbb{C} \times T_f\mathrm{Map}(S^2, M) \otimes \mathbb{C} \longrightarrow \mathbb{C}$$

given by

$$d^2E(f)(V, \bar{V}) = 4\int_{S^2}\left[\left|\frac{DV}{\partial \bar{z}}\right|^2 - \left\langle \mathcal{R}\left(V \wedge \frac{\partial f}{\partial z}\right), \bar{V} \wedge \frac{\partial f}{\partial \bar{z}}\right\rangle\right] dxdy, \tag{4}$$

in which

$$\frac{\partial f}{\partial z} = \frac{1}{2}\left(\frac{\partial f}{\partial x} - i\frac{\partial f}{\partial y}\right) \quad \text{and} \quad \frac{D}{\partial \bar{z}} = \frac{1}{2}\left(\frac{D}{\partial \bar{x}} + i\frac{D}{\partial \bar{y}}\right)$$

is the Cauchy-Riemann operator defined by the Levi-Civita connection on $\mathbf{E} = f^*TM \otimes \mathbb{C}$. If σ is the two-plane spanned by $\partial f/\partial z$ and V, then (5) can also be written as

$$d^2E(f)(V, \bar{V}) = 2\int_{S^2}\left[\left\|\bar{\partial} V\right\|^2 - K(\sigma)e(f)\left\|V\right\|^2\right] dA, \tag{5}$$

where $K(\sigma)$ is the complex sectional curvature of σ, $e(f)$ is the energy density of f and dA is the element of area on S^2 when the S^2 is given the canonical Riemannian metric of constant curvature and total area one.

The Riemann-Roch Theorem (see [22] or [29]) allows us to estimate the dimension of the space of solutions to the equation $\bar{\partial} V = 0$, and if the complex sectional curvature is positive, this enables us to give a lower bound on the *Morse index* of f, which by definition is the maximum dimension of a complex linear space of sections of $\mathbf{E} = f^*TM \otimes \mathbb{C}$ on which $d^2E(f)$ is negative definite.

Definition. A two plane $\sigma \subseteq T_pM \otimes \mathbb{C}$ is *isotropic* if it is spanned by linearly independent vectors v and w such that

$$\langle v, v\rangle = \langle v, w\rangle = \langle w, w\rangle = 0.$$

It is *half-isotropic* if it is spanned by linearly independent vectors v and w such that

$$\langle v, v\rangle = \langle v, w\rangle = 0.$$

Since $v = x + iy$ is isotropic, x and y are orthogonal and thus span a real two plane $\widehat{\sigma}$ which we call a real two-plane *associated* to the complex two-plane σ.

Half-isotropic curvature was discussed briefly in §5 of [22]. Note that positive half-isotropic curvature reduces to positive Ricci curvature when M has dimension three, and the two-planes which appear in the second variation formula (5) are half-isotropic. It is the sectional curvature of half-isotropic two-planes which governs stability of minimal two-spheres, just as the real sectional curvatures govern stability of geodesics.

Lemma 1.2. *Pinching of the real sectional curvatures $K_r(\sigma_0)$ and the half-isotropic curvatures $K_i(\sigma_1)$ are related by*

$$\delta < K_r(\sigma_0) \leq 1 \quad \Rightarrow \quad \frac{4}{3}\delta - \frac{1}{3} < K_i(\sigma_1) \leq \frac{4}{3} - \frac{\delta}{3}. \tag{6}$$

Indeed if (e_1, e_2, e_3, e_4) are unit-length vectors, perpendicular to each other, at a point p, and σ_1 is the half-isotropic two-plane spanned by $e_1 + ie_2$ and $ae_3 + ibe_4$, where $a, b \in \mathbb{R}^+$, then

$$\begin{aligned}\langle \mathcal{R}((e_1 + ie_2) \wedge (ae_3 + ibe_4)), (e_1 - ie_2) \wedge (ae_3 - ibe_4)\rangle \\ = a^2(R_{1313} + R_{2323}) + b^2(R_{1414} + R_{2424}) + 2abR_{1243},\end{aligned}$$

and according to Berger's formula (7) in [3], $|R_{1243}| \leq (2/3)(1 - \delta)$, so

$$\begin{aligned}&\delta < K_r(\sigma_0) \leq 1 \quad \Rightarrow \\ &\quad 2\delta(a^2 + b^2) - \frac{4}{3}ab(1 - \delta) < 2(a^2 + b^2)K_i(\sigma_1) \leq 2(a^2 + b^2) + \frac{4}{3}ab(1 - \delta),\end{aligned}$$

which simplifies to the conclusion of the lemma:

$$\delta - \frac{1}{3}(1 - \delta) < K_i(\sigma_1) \leq 1 + \frac{1}{3}(1 - \delta).$$

The Lemma shows that if the real sectional curvatures are 1/4-pinched, the half-isotropic curvatures are positive, and that the hypothesis (2) implies that

$$\frac{1}{3} < K_i \quad \text{and} \quad K_r \leq 1. \tag{7}$$

It is actually this weaker hypothesis that is needed in the proof of Theorem 1.1.

Unlike the case of geodesics, the Morse index of minimal two-spheres depends only on pointwise pinching, and the curvature hypothesis in Theorem 1.1 can be weakened: one only needs a relation between the complex sectional curvature

of a half-isotropic two-plane σ and the real sectional curvature of an associated real two-plane $\widehat{\sigma}$.

Theorem 1.3. *Suppose that the compact smooth simply connected manifold M has dimension $n \geq 4$, and has a Riemannian metric which satisfies the condition*

$$K_i(\sigma) > \frac{1}{3} K_r(\widehat{\sigma}) > 0, \tag{8}$$

where $\widehat{\sigma}$ is a real two-plane associated to σ. If all prime minimal two-spheres of Morse index $\leq 2n-5$ in M are immersions and lie on Morse nondegenerate critical submanifolds, each a $\widehat{\mathcal{G}}$-orbit, then the number of distinct prime minimal two spheres of Morse index λ, none of which double cover minimal projective planes, is at least $p_3(\lambda - n + 2)$ for $n - 2 \leq \lambda \leq 2n - 5$, where $p_3(k)$ is the number of k-cells in the Schubert cell decomposition for $G_3(\mathbb{R}^{n+1})$.

We remark that the condition on prime minimal two-spheres is satisfied for generic metrics by the Bumpy Metric Theorem presented in [29]. We will see that it follows from the second variation formula (5) that any minimal two-sphere $f : S^2 \to M$ in a manifold of positive complex sectional curvatures must have Morse index at least $n - 2$, where n is the dimension of M. We will later use this fact to prove the following step towards the above theorems:

Nonbubbling Theorem 1.4. *Suppose that the n-dimensional Riemannian manifold M has positive complex sectional curvature, $n \geq 4$. If a sequence of α-energy critical points approaches a bubble tree containing k nonconstant minimal two-spheres in the limit, then the Morse index of these critical points must eventually be at least $k(n-2)$.*

To prove Theorem 1.3, we use the Nonbubbling Theorem to control bubbling, the first of the two problems we mentioned before and use the curvature hypothesis (7) to show that all branched covers of order $d \geq 3$ must have Morse index at least $2n - 4$. This allows us to limit consideration to only two-fold branched covers, and finally we use equivariant Morse theory with $\mathbb{Z}_2$ coefficients to rule out branched covers of order two, as well as double covers of projective planes. Our goal was to develop techniques that will be useful in related problems, such as minimal surfaces of low index in symmetric spaces of compact type sufficiently close to standard metrics.

In § 2, we describe a center of mass technique which allow us to reduce the noncompact symmetry group $\widehat{\mathcal{G}}$ for E to the compact symmetry group $\widehat{G} = O(3)$ needed for E_α, and in the following section we show that critical points for E_α always have finite isotropy. In § 4 we explain how to extend Uhlenbeck's Morse theory on Banach manifolds to the G-equivariant case. We next give conditions in § 5 under which the number of critical points of index less than a given bound is finite, a weak version the compactness theorem of Choi and Schoen [8] for imbedded minimal surfaces in three-manifolds. This compactness theorem was used by White [40] and [39] to prove existence of at least one minimal torus in a three-sphere with positive Ricci curvature, suggesting the relevance

of positive half-isotropic curvature in our case, which reduces to positive Ricci curvature for three-manifolds. We then prove the Nonbubbling Theorem in § 6. We study the space of d-fold covers of a prime imbedding in §7, providing an index estimate which rules out the case $d \geq 3$ and describing an explicit model by Milnor [25] for treating the case $d = 2$. We then review equivariant topology and describe its calculation for the space of maps $f : S^2 \to S^n$ of energy strictly less than 8π in §8. The Morse-Witten chain complex approach to Morse theory is described in §9, and we show that G-equivariant Morse functions are dense by an argument of Wassermann [38] extended to infinite-dimensions; see also [1]. We then have the ingredients necessary to reduce the proof of Theorems 1.1 and 1.3 to a calculation in equivariant cohomology following Bott [4] and Hingston [16]. The fact that we need consider only two-fold covers allows us to use $\mathbb{Z}_2$-coefficients which make the calculations more tractible.

The techniques developed here are improvements on those used in [28]. Some readers may find it useful to consult [29] which gives background for this article.

2 Perturbations of minimal surfaces

Our strategy is to show that a harmonic two-sphere in M can be perturbed to a family of α-harmonic two-spheres, depending on the parameter α, for $\alpha \in (1, \alpha_0)$ for some $\alpha_0 > 1$, but we must deal with the technical problem that the group of symmetries changes when the usual energy E is perturbed to the α-energy E_α for $\alpha > 1$. While the usual energy E is invariant under the group

$$\widehat{\mathcal{G}} = PSL(2, \mathbb{C}) \cup R \cdot PSL(2, \mathbb{C}), \quad R \text{ a reflection},$$

of conformal and anticonformal transformations of the Riemann sphere, when $\alpha > 1$, the α-energy

$$E_\alpha : C^2(S^2, M) \to \mathbb{R} \quad \text{defined by} \quad E_{S^2}(f) = \frac{1}{2} \int_\Sigma (1 + |df|^2)^\alpha dA - \frac{1}{2}$$

is invariant only under the smaller group $O(3)$ of isometries. This smaller group of symmetries is reflected in an additional condition on critical points of E_α: they must be parametrized so that a certain "center of mass" is zero:

Lemma 2.1. *Let $\mathbf{X} : S^2 \to \mathbb{R}^3$ denote the standard inclusion, and let $\mathbf{0}$ denote the origin in $\mathbb{R}^3$. For each $\alpha \in (1, \infty)$, there is a smooth function $\psi_\alpha : [0, \infty) \to \mathbb{R}$ such that $\psi_\alpha(0) = 0$, $\psi_\alpha(t), \psi'_\alpha(t) > 0$ for $t > 0$, and if f is a critical point for E_α, then*

$$\int_{S^2} \mathbf{X} \psi_\alpha(|df|^2) dA = \mathbf{0}. \tag{9}$$

In the argument for this lemma, we regard S^2 as the one-point compactification $\mathbb{C} \cup \{\infty\}$ of the complex plane $\mathbb{C}$ with the standard coordinate

$$z = x + iy = re^{i\theta} = e^{u+i\theta}, \tag{10}$$

and give S^2 the constant curvature metric expressed in polar coordinates by

$$g = ds^2 = \frac{4}{(1+r^2)^2}(dr^2 + r^2 d\theta^2) = \frac{1}{\cosh^2 u}(du^2 + d\theta^2), \tag{11}$$

where $r = e^u$. (This metric is related by scaling to the constant curvature metric of total area one.) For $t \in \mathbb{R}$, we define a family of linear fractional transformations,

$$\phi_t : S^2 \to S^2 \quad \text{by} \quad u \circ \phi_t = u + t, \quad \theta \circ \phi_t = \theta,$$

so that $t \mapsto \phi_t$ is a one-parameter subgroup of conformal transformations. The energy density is then given by the formula

$$e(f \circ \phi_t) = \frac{1}{2}\left(\left|\frac{\partial f}{\partial u}\right|^2 + \left|\frac{\partial f}{\partial \theta}\right|^2\right)^2 \cosh^2(u+t),$$

and it is straightforward to calculate its derivative at $t = 0$:

$$\begin{aligned}\frac{d}{dt} e(f \circ \phi_t)\bigg|_{t=0} &= \frac{1}{2}\,\frac{d}{dt}|d(f \circ \phi_t)|^2\bigg|_{t=0} \\ &= \left(\left|\frac{\partial f}{\partial u}\right|^2 + \left|\frac{\partial f}{\partial \theta}\right|^2\right) \sinh u \cosh u = |df|^2 \tanh u. \end{aligned} \tag{12}$$

In terms of the standard Euclidean coordinates (x, y, z) on $\mathbb{R}^3$, S^2 is represented by the equation $x^2 + y^2 + z^2 = 1$, and a straightforward computation using stereographic projection from the north pole to the (x, y)-plane shows that

$$z = \frac{r^2 - 1}{r^2 + 1} = \frac{\sinh u}{\cosh u} = \tanh u.$$

We can use (12) to calculate the derivative of E_α, and obtain

$$\begin{aligned}\frac{d}{dt} E_\alpha(f \circ \phi_t)\bigg|_{t=0} &= \frac{1}{2}\,\frac{d}{dt}\int_{S^2} (1 + |d(f \circ \phi_t)|^2)^\alpha \operatorname{sech}^2(u+t)\, du d\theta\bigg|_{t=0} \\ &= \alpha \int_{S^2} (1 + |df|^2)^{\alpha-1} \tanh u |df|^2 dA - \int_{S^2} (1 + |df|^2)^\alpha \tanh u \; dA.\end{aligned}$$

Thus, if f is a critical point for E_α,

$$\begin{aligned} 0 &= \int_{S^2} [\alpha(1 + |df|^2)^{\alpha-1}|df|^2 - (1 + |df|^2)^\alpha] z dA, \\ &= \int_{S^2} [\alpha(1 + |df|^2)^{\alpha-1}|df|^2 - (1 + |df|^2)^\alpha + 1] z dA, \end{aligned}$$

where we have used the fact that the average value of z on S^2 is zero. We obtain (9) by setting

$$\psi_\alpha(t) = \frac{\alpha(1+t)^{\alpha-1}t - (1+t)^\alpha + 1}{\alpha - 1} = \int_0^t \alpha \frac{\tau}{(1+\tau)^{2-\alpha}} d\tau. \tag{13}$$

One easily verifies that this function has the desired properties, and has a smooth limit as $\alpha \to 1$, namely

$$\psi_1(t) = \int_0^t \frac{\tau}{(1+\tau)} d\tau = t - \log(1+t). \tag{14}$$

Definition. We will say that a smooth map $f : S^2 \to M$ has *α-center of mass zero* if it satisfies the condition

$$\int_{S^2} \mathbf{X} \psi_\alpha(|df|^2) dA = \mathbf{0}, \tag{15}$$

where $\mathbf{X} : S^2 \to \mathbb{R}^3$ is the standard inclusion as a round sphere centered at the origin. and we let

$$\mathrm{Map}_0(S^2, M) = \left\{ f \in \mathrm{Map}(S^2, M) : \int_{S^2} \mathbf{X} \psi_1(|df|^2) dA = \mathbf{0} \right\}.$$

Note that if $T \in \widehat{\mathcal{G}}$,

$$f,\ f \circ T \in \mathrm{Map}_0(S^2, M) \quad \text{and } f \text{ nonconstant} \quad \Rightarrow \quad T \in O(3),$$

so setting the center of mass zero does indeed cut the symmetry group down to $O(3)$.

Theorem 2.2. *Suppose that M is a compact connected Riemannian manifold and that $f : S^2 \to M$ is a prime oriented parametrized minimal immersion lying on a nondegenerate critical submanifold N for E of dimension six, an orbit for the $PSL(2, \mathbb{C})$ action described above. Let $N_0 = N \cap Map_0(S^2, M)$. Then for some $\alpha_0 \in (1, \infty)$, there is a smooth map*

$$\eta : N_0 \times [1, \alpha_0) \longrightarrow \left\{ (f, \alpha) \in L^2_k(S^2, M) \times [1, \alpha_0) : \int_{S^2} \mathbf{X} \psi_\alpha(|df|^2) dA = \mathbf{0} \right\}$$

which satisfies the following conditions:

1. *$\eta(f, \alpha)$ is a critical point for E_α for each $f \in N_0$ and $\alpha \in [1, \alpha_0)$,*
2. *η is $O(3)$-equivariant, and*
3. *there is an $\epsilon_0 > 0$ such that if f' is a critical point for E_α such that $\|f' - f\|_{C^2} < \epsilon_0$ for some $f \in N_0$, f' lies in the image of η.*

Note that it follows from Condition 2 that if $f \circ A = f$ for $f \in N_0$, where $A : S^2 \to S^2$ is the antipodal map, then

$$\eta(f, \alpha) \circ A = \eta(f, \alpha), \quad \text{for } f \in N_0 \text{ and } \alpha \in [1, \alpha_0).$$

Once we have dealt with the change in symmetry group, the proof of Theorem 2.2 is a relatively direct consequence of the implicit function theorem. Note that when k is relatively large, the map

$$E_\alpha : L^2_k(S^2, M) \longrightarrow \mathbb{R}$$

is actually C^∞; indeed, we can regard E_α as a composition of several maps

$$f \mapsto df \mapsto (1 + |df|^2)^\alpha \mapsto \int_\Sigma (1 + |df|^2)^\alpha dA,$$

each of which is smooth. The first map is smooth into the Hilbert manifold $L^2_{k-1}(S^2, TM)$, the second is smooth by the well-known ω-Lemma, while the third map is always smooth since integration is continuous and linear. Similarly, the two-variable map

$$E^\star : L^2_k(S^2, M) \times [1, \alpha_0) \longrightarrow \mathbb{R} \quad \text{defined by} \quad E^\star(f, \alpha) = E_\alpha(f),$$

is also smooth when $\alpha_0 > 1$, as is its restriction to

$$\mathcal{M}_0 = \left\{ (f, \alpha) \in L^2_k(S^2, M) \times [1, \alpha_0) : \int_{S^2} \mathbf{X}\psi_\alpha(|df|^2) dA = \mathbf{0} \right\}.$$

Of course, when $\alpha = 1$ we obtain the usual energy, $E^\star(f, 1) = E(f)$ restricted to the L^2_k-completion of $\mathrm{Map}_0(S^2, M)$.

Choose $f_0 \in N_0$ and let S_{f_0} be a slice of the $SO(3)$-action through $(f_0, 1) \in \mathcal{M}_0$. We choose a coordinate system

$$\phi : S_{f_0} \longrightarrow V \times [1, \alpha_0) \subseteq \mathbf{H} \times [1, \alpha_0),$$

where V is a convex open neighborhood of 0 in a Hilbert space $\mathbf{H}$ with inner product $\langle \cdot, \cdot \rangle$, such that

$$\phi(f, \alpha) = (\phi_0(f, \alpha), \alpha).$$

Using this coordinate system we create a corresponding smooth function

$$F^\star : V \times [1, \alpha_0) \longrightarrow \mathbb{R}, \quad F^\star(\phi_0(f, \alpha), \alpha) = E^\star(f, \alpha).$$

We then define a vector field $\mathcal{X} : V \times [1, \alpha_0) \to \mathbf{H}$ by

$$D_1 F^\star(x, \alpha) = \langle \mathcal{X}(x), y \rangle, \quad \text{for } x, y \in \mathbf{H}.$$

Since the null space for $d^2E(f)$ is generated by the symmetry group G, the restriction of $d^2E(f)$ to S_{f_0} is nondegenerate, and this implies that $D_1\mathcal{X}(0)$, the partial derivative with $\alpha = 1$ held constant, is invertible. The implicit function theorem (eg Theorem 5.9 in Chapter I of [19]) implies that there is a smooth map $g : [1, \alpha_0) \to \mathbf{H}$ with $g(1) = 0$ such that

$$\mathcal{X}(g(\alpha), \alpha) = 0.$$

This in turn yields a smooth curve $\tilde{g} : [1, \alpha_0) \to \mathcal{M}_0$, $\tilde{g}(\alpha) = (\eta(\alpha), \alpha)$, in which each $\eta(\alpha)$ is a critical point for the restriction of E_α to the space of maps satisfying (15). But our calculation at the beginning of the section shows that these are also critical for conformal dilations, hence critical for all variations within $L^2_k(S^2, M)$.

Thus, after possibly decreasing $\alpha_0 > 1$, we obtain a smooth curve of critical points

$$\alpha \mapsto \eta(\alpha) = \eta(f_0, \alpha)$$

for E_α. We then extend η to $N_0 \times [1, \alpha_0)$ by demanding $SO(3)$-equivariance, and the resulting extension satisfies all the conditions of Theorem 3.2.

3 Isotropy groups

When developing general theory for the next few sections, we assume that M is a compact smooth manifold with an arbitrary Riemannian metric. The α-energy

$$E_\alpha : \mathrm{Map}(S^2, M) \longrightarrow \mathbb{R}$$

is invariant under an action of the group $\widehat{G} = O(3)$ by isometries on the domain. Because of this action, the nonconstant critical points for E_α lie on $\widehat{G}$-orbits of critical points, and our goal is to apply $\widehat{G}$-equivariant Morse theory, in which isotropy groups of the $\widehat{G}$-action play an important role.

The isotropy group of any point in $\mathrm{Map}(S^2, M)$ must be a closed subgroup of $\widehat{G}$, hence a Lie subgroup. One possibility it that the isotropy group is $\widehat{G}$ itself, and this occurs exactly for the constant maps, which form a submanifold M_0 of zero energy which is diffeomorphic to M itself. If the isotropy group $H \subseteq \widehat{G}$ of a nonconstant map is not discrete, the only possibility for its identity component is S^1, a group of rotations about some axis.

If the isotropy group contains S^1, a smooth map $f : S^2 \to M$ must in fact degenerate to a curve parametrized by S^2, so we examine whether this could happen for a critical point for E_α. As in the previous section, we give S^2 the constant curvature metric expressed in polar coordinates by

$$g = ds^2 = \frac{4}{(1+r^2)^2}(dr^2 + r^2 d\theta^2) = \frac{1}{\cosh^2 u}(du^2 + d\theta^2),$$

where $r = e^u$, chosen so that a rotation through an angle ϕ is represented by $\theta \mapsto \theta + \phi$. Then critical points for E_α which are invariant under S^1 are simply critical points independent of θ, so we restrict E_α to functions $f_0(u)$ of the variable u. Then E_α simplifies to

$$E_\alpha(f_0) = \pi \int_{-\infty}^{\infty} \left[1 + (\cosh u \|f_0'(u)\|)^2\right]^\alpha \operatorname{sech}^2 u \; du - \frac{1}{2},$$

where $f_0 : \mathbb{R} \to M$ is now a smooth curve, with tangent vector $f_0'(u)$, at $f_0(u)$, which has length $\|f_0'(u)\|$ in terms of the Riemannian metric on M. For f_0 to

be a critical point, it must satisfy the Euler-Lagrange equation, which is simply

$$\frac{D^g}{du}\left\{\left[1+(\cosh u\|f_0'(u)\|)^2\right]^{\alpha-1} f_0'(u)\right\}=0,$$

where D^g is the covariant derivative defined by the Riemannian metric on M, and this equation has as solutions $f_0=\gamma\circ h$, where $\gamma:\mathbb{R}\to M$ is a unit speed geodesic and $h:\mathbb{R}\to\mathbb{R}$ is a smooth function such that

$$\left[1+(\cosh u\ h'(u))^2\right]^{\alpha-1} h'(u)=a,$$

with a being a constant. An important feature of this equation is that $h'(u)$ assumes its minimum value when $u=0$, and hence

$$h'(u)\geq c \quad \text{for all } u, \quad \text{where} \quad g(c)=\left[1+c^2\right]^{\alpha-1} c=a.$$

Since g is a strictly increasing function, there is a unique positive solution c to the equation $g(c)=a$, and

$$\begin{aligned} E_\alpha(f_0)&\geq \pi\int_{-\infty}^{\infty}\left[1+c^2\cosh^2 u\right]^{\alpha}\operatorname{sech}^2 u\ du-\frac{1}{2}\\ &\geq \pi\int_{-\infty}^{\infty} c^{2\alpha}\cosh^{2(\alpha-1)} u\ du-\frac{1}{2}=\infty. \end{aligned}$$

Thus we see that there are no nonconstant finite energy critical points to E_α which have isotropy group containing S^1, or:

Lemma 3.1. *Critical points for the α-energy $E_\alpha:Map(S^2,M)\to\mathbb{R}$ of finite α-energy are either constant with isotropy group $\widehat{G}=O(3)$, or their isotropy groups for the $\widehat{G}$-action are finite.*

Of course, $\mathrm{Map}(S^2,M)-M_0$ contains many points at which the isotropy group of the $\widehat{G}$-action is a circle, but none of these are critical points for E_α. For a given choice of Riemannian metric on M, we let K denote the set of critical points for

$$E_\alpha:\mathrm{Map}(S^2,M)-M_0\longrightarrow\mathbb{R}.$$

Then Lemma 3.1 implies that the action of $\widehat{G}=O(3)$ on an open neighborhood of K is locally free. Since the local flow for the gradient of α-energy preserves isotropy groups, it is also the case that the $\widehat{G}$-action is locally free on a neighborhood U of all trajectories for this gradient flow which connect points of K.

We can also ask what happens when the action of the smaller group $G=SO(3)$ fails to be free at a critical point $f:S^2\to M$ for either E or E_α. By Lemma 2.1, the isotropy group must be finite, and the finite subgroups of $SO(3)$ are classified in §2.6 of [41]. The only possibilities are cyclic groups, dihedral groups and symmetry groups for the five platonic solids. For each of these groups, there are points on S^2 with nontrivial isotropy, and in the case of harmonic maps, such points must be branch points.

Under the assumption that all prime minimal two-spheres of Morse index $\leq 2n-5$ are free of branch points, the isotropy of the $SO(3)$ action at these critical points must be trivial. Critical points for E_α which closely approximate these minimal two-spheres must also have trivial $SO(3)$ isotropy.

4 Morse theory on Banach manifolds

Suppose that $\mathcal{M}$ is a smooth manifold (say an infinite-dimensional Banach manifold) with a continuous right G-action by diffeomorphisms, where G is a compact Lie group. If $F : \mathcal{M} \to \mathbb{R}$ is a G-invariant C^2 function, then F descends to a map $F_0 : \mathcal{M}/G \to \mathbb{R}$, where $\mathcal{M}/G$ is the space of orbits. When the action of G is free, the quotient is a well-behaved topological manifold. If the action is only "locally free," that is the isotropy groups are finite, we can apply the slice and tube theorems, extended from §2.3 and 2.4 of [9] to Banach manifolds, and study F by means of its restriction to a slice $\mathcal{S}_{f_0}$ through a given point $f_0 \in \mathcal{M}$.

There are three approaches to constructing a Morse theory for such a function F:

1. We can perturb F to a *Morse function* which has nondegenerate critical points in the usual sense, but this breaks the symmetry under the G-action.

2. We can look for a perturbation of F to an *equivariant Morse function* for which the critical points lie on nondegenerate critical submanifolds (as defined by Bott), each of which is a single G-orbit. If the action is locally free, the restriction of F to any slice has critical points which are nondegenerate in the usual sense.

3. We can seek a a *Morse-Bott function* on $\mathcal{M}$ for which all critical points lie on finite-dimensional compact nondegenerate critical submanifolds (which may be larger than a single G-orbit).

Each of the viewpoints has advantages and disadvantages. We are interested in applying the second and third cases to the α-energy $E_\alpha : \mathcal{M} \to [0, \infty)$, where $\mathcal{M} = L_1^{2\alpha}(S^2, M)$, M being a compact Riemannian manifold. On the other hand, the first case can be applied to a suitable perturbation of the α-energy, say $F : \mathcal{M} \to \mathbb{R}$, where

$$F_\alpha(f) = \frac{1}{2}\int_{S^2}(1+|df|^2)^\alpha dA - \frac{1}{2} + \int_{S^2} f \cdot \psi dA, \tag{16}$$

the last perturbation term being defined in terms of an isometric immersion of M into some Euclidean space $\mathbb{R}^N$ and via a suitable small function $\psi : S^2 \to \mathbb{R}^N$. In all cases the function is C^2, satisfies Condition C and has C^∞ critical points. It is proven in [22] or [29] that for generic choices of ψ, the function F defined by (16) is bounded below and has nondegenerate critical points.

The first step in constructing a Morse theory consists of determining formulae for the second variation of E_α or F. A calculation (carried out in [22] or

[29]) shows that the second variation of E_α at a critical point $f_0 \in \mathcal{M}$ is the symmetric bilinear form

$$d^2E_\alpha(f_0) : T_{f_0}L_1^{2\alpha}(S^2, M) \times T_{f_0}L_1^{2\alpha}(S^2, M) \longrightarrow \mathbb{R}$$

defined by

$$\begin{aligned} d^2E_\alpha(f_0)(X, Y) = \alpha \int_{S^2} (1 + |df_0|^2)^{\alpha-1}[\langle DX, DY\rangle - \langle \mathcal{K}(X), Y\rangle]dA \\ + 2\alpha(\alpha - 1) \int_{S^2} (1 + |df_0|^2)^{\alpha-2}\langle df_0, DX\rangle\langle df_0, DY\rangle dA, \quad (17) \end{aligned}$$

where D is the covariant differential and the bundle endomorphism $\mathcal{K}$ is determined at p in terms of an orthonormal basis (e_1, e_2) for T_pM by

$$\langle \mathcal{K}(X), Y\rangle(p) = \sum_{i=1}^{2} \langle \mathcal{R}(X(p) \wedge e_i), Y(p) \wedge e_i\rangle \,.$$

This second variation (17) approaches the second variation (5) for the usual energy E as $\alpha \to 1$. Similarly, the second variation of F is given by

$$\begin{aligned} d^2F(f_0)(X, Y) = \alpha \int_{S^2} (1 + |df_0|^2)^{\alpha-1}[\langle DX, DY\rangle - \langle \mathcal{K}(X), Y\rangle]dA \\ + 2\alpha(\alpha - 1) \int_{S^2} (1 + |df_0|^2)^{\alpha-2}\langle df_0, DX\rangle\langle df_0, DY\rangle dA + \int_{S^2} \psi \cdot \alpha(X, Y)dA, \end{aligned} \quad (18)$$

where α is the second fundamental form of M in $\mathbb{R}^N$. We can define the *Morse index* of an element $f_0 \in L_1^{2\alpha}(S^2, M)$, whether f_0 is critical or not, to be the maximal dimension of a linear subspace of

$$T_{f_0}L_1^{2\alpha}(S^2, M)$$

on which d^2F is negative-definite; note that the Morse index is a lower semi-continuous function of $f_0 \in \mathcal{M}$. It will be useful to compare these formulae to an inner product $\langle\langle \cdot, \cdot \rangle\rangle$ we can define by

$$\langle\langle X, Y\rangle\rangle = 2\alpha \int_{S^2} (1 + |df_0|^2)^{\alpha-1}[\langle DX, DY\rangle + \langle X, Y\rangle]dA, \quad (19)$$

which is equivalent to the usual L_1^2 inner product, when f_0 is a fixed C^∞ map.

We now review the Morse theory on Banach manifolds of Uhlenbeck [37], together with its straightforward extension to equivariant Morse functions or Morse-Bott functions. For this, it is convenient to extend the notion of "gradient-like" vector field $\mathcal{X}$ (as described in Milnor [24]) to the three cases. In each case, the gradient-like vector field $\mathcal{X}$ is constructed by an explicit formula near the critical locus, which is then pieced together with a pseudogradient as described by Palais [30] away from the critical locus.

I. Near the critical locus. Consider first the function $F : \mathcal{M} \to [0, \infty)$ defined by (16) which is Morse nondegenerate in the usual sense; the critical points of a such a function are isolated. Our first requirement is that the restriction of $\mathcal{X}$ to an open neighborhood $U_{f_0} = U$ of a critical point $f_0 \in \mathcal{M}$ is replaced by a "canonical Fredholm map" A which is transported via an exponential chart to U,

$$A : T_{f_0}\mathcal{M} \longrightarrow T_{f_0}\mathcal{M} \quad \mapsto \quad \mathcal{X}|U = \mathcal{X}_A = \exp_{f_0} \circ A \circ (\exp_{f_0})^{-1}|U. \tag{20}$$

This canonical Fredholm map A is defined in terms of the second variation (18) and the inner product (19) by

$$d^2F(f_0)(A(X), Y) = \langle\langle X, Y \rangle\rangle. \tag{21}$$

(Thus A^{-1} would define a gradient with respect to the inner product.) It is straightforward to see that A is a bounded symmetric endomorphism in terms of the L^2_1 norm and that it is also bounded in terms of the L^2_k norms for $k \in \mathbb{N}$. Lemma 5.3 of [37] shows that A restricts to a bounded endomorphism of $L^{2\alpha}_1(S^2, f_0^*TM)$, while other estimates in [37] show that $\mathcal{X}_A$ satisfies the condition

$$dF(\mathcal{X}_A)(p) > \varepsilon_0 \|p\|^2 > 0 \quad \text{for } p \in U - \{f_0\}, \tag{22}$$

for some $\varepsilon_0 > 0$, where $\| \cdot \|$ denotes the $L^{2\alpha}_1$ norm in exponential coordinates. This is the condition used by Uhlenbeck [37] for the definition of "weak gradient," and is sufficient for constructing a handle-body decomposition.

For the other cases, cases in which

$$E_\alpha : \mathcal{M} \to \mathbb{R}, \quad \mathcal{M} = L^{2\alpha}_1(S^2, M)$$

is equivariant Morse or Morse-Bott, the critical locus is the union of compact nondegenerate critical submanifolds. If K is one of those nondegenerate critical submanifolds and $f_0 \in K$, we can use the inner product (19) to divide $T_{f_0}\mathcal{M}$ into an orthogonal direct sum decomposition,

$$T_{f_0}\mathcal{M} = T_{f_0}K \oplus S_{f_0},$$

with $T_{f_0}K$ being not only the tangent space to K, but also the null space of the symmetric bilinear form (17). As f_0 varies through K, we get a direct sum decomposition

$$T\mathcal{M}|K \oplus S,$$

where S is the normal bundle to K in $\mathcal{M}$. For any $f_0 \in K$, we can then define $A_{f_0} : S_{f_0} \to S_{f_0}$ by (21), a linear map which is bounded with respect to the $L^{2\alpha}_1$ norm or any of the L^2_k norms restricted to S_{f_0}.

The family of maps $f_0 \to A_{f_0}$ defines a vector field $\tilde{\mathcal{X}}_A$ on the total space of S which is tangent to the fibers. Finally, there is a diffeomorphism ϕ_K from a neighborhood V of the zero section in this normal bundle to an open

neighborhood $U = U_K$ of K such that ϕ_K and $(\phi_K)_*$ are the identity along the zero section, and we set $\mathcal{X}_A = (\phi_K)_*(\tilde{\mathcal{X}}_A)$. This field satisfies

$$dF(\mathcal{X}_A)(p) > \varepsilon_0 \|v\|^2 > 0 \quad \text{for } p = \phi_K(v), v \in S_{f_0}, \tag{23}$$

for some $\varepsilon_0 > 0$, where $\|\cdot\|$ denotes the $L_1^{2\alpha}$ norm on S_{f_0}.

Note that the Morse-Bott case contains the others as special cases.

II. Away from the critical locus. The second requirement is that away from the critical locus, the vector field is a *pseudogradient* as described by Palais [30]; that is, there exist constants $\varepsilon_1 > 0$ and $\varepsilon_2 > 0$ such that

$$\|\mathcal{X}(p)\| < \varepsilon_1 \|dF_p\|, \quad \|dF_p\|^2 < \varepsilon_2 dF_p(\mathcal{X}(p)). \tag{24}$$

It is important to note that when these estimates hold for $\mathcal{X}_0$ and $\mathcal{X}_1$, they hold for any convex combination thereof, and therefore pseudogradients can be pieced together by the C^2 partitions of unity available on the Banach manifold $L_1^{2\alpha}(S^2, f_0^*TM)$. These estimates imply a third,

$$\|\mathcal{X}(p)\|^2 < \varepsilon_1^2 \varepsilon_2 dF_p(X(p)). \tag{25}$$

which in turn implies that the flow for $-\mathcal{X}$ will be positively complete by the following argument:

Let $\{\phi_t : t \in \mathbb{R}\}$ denote the local one-parameter group generated by $-\mathcal{X}$, and let $d : \mathcal{M} \times \mathcal{M} \to \mathbb{R}$ denote the distance function defined by the Finsler metric which $\mathcal{M}$ inherits from the ambient Finsler manifold $L_1^{2\alpha}(S^2, \mathbb{R}^N)$. Then it follows from (25) and the Cauchy-Schwarz inequality that if $t_1 < t_2$,

$$\begin{aligned} d(\phi_{t_1}(p), \phi_{t_2}(p))^2 &\le \left[\int_{t_1}^{t_2} \left\| \frac{d}{dt}\phi_t(p) \right\| dt\right]^2 = \left[\int_{t_1}^{t_2} \|-\mathcal{X}(\phi_t(p))\|\, dt\right]^2 \\ &\le (t_2 - t_1)\left[\int_{t_1}^{t_2} \|\mathcal{X}(\phi_t(p))\|^2 dt\right] \le \varepsilon_1^2 \varepsilon_2 (t_2 - t_1) \int_{t_1}^{t_2} dF(\mathcal{X})(\phi_t(p)) dt \\ &= \varepsilon_1^2 \varepsilon_2 (t_2 - t_1)(F(\phi_{t_1}(p)) - F(\phi_{t_2}(p))). \end{aligned} \tag{26}$$

Since F is bounded below, this inequality implies that if $\phi_t(p)$ is defined only for $t \in [0, b)$, for some finite b, then any sequence $t_i \to b$ yields a Cauchy sequence $\{\phi_{t_i}(p)\}$, and completeness of the Finsler metric shows that this Cauchy sequence must converge to some $q \in \mathcal{M}$. But there is a family of integral curves passing through points near q, showing that $[0, b)$ is not a maximal interval on which $\phi_t(p)$ is defined.

On the other hand, the second inequality of (24) shows that $\|dF\| \to 0$ along a sequence of points on each orbit, and then Condition C implies that each orbit eventually enters a neighborhood of some critical point. Since the function E_α or F satisfies Condition C, each component of the critical set K will be compact.

Definition. Suppose that $F : \mathcal{M} \to \mathbb{R}$ is a C^2 Morse-Bott function on the Banach manifold $\mathcal{M}$ which satisfies Condition C and is bounded below. A

gradient-like vector field for F is a vector field $\mathcal{X}$ on $\mathcal{M}$ which satisfies the following requirements: If K_0 is a component of the critical locus and thus a compact nondegenerate critical submanifold, there exist open neighborhoods

$$V(K_0) = \bigcup\{V_{f_0} : f_0 \in K_0\} \quad \text{and} \quad U(K_0) = \bigcup\{U_{f_0} : f_0 \in K_0\}$$

where V_{f_0} is an open ball neighborhood of f_0 in terms of exponential coordinates, with $\overline{V(K_0)} \subseteq U(K_0)$ such that $\mathcal{X}$ satisfies (20) and (22) on U_{f_0} for each $f_0 \in K_0$. Moreover, $\mathcal{X}$ is a pseudogradient on

$$\mathcal{M} - \bigcup\{\overline{V(K_0)} : \ K_0 \text{ a component of the critical locus } \}.$$

A gradient-like vector field is also positively complete by the same argument we gave for pseudogradients, except we replace estimate (25) by a combination of estimates (22) and (25). By Condition C, there are only finitely many critical submanifolds with values of F no larger than a given energy bound E_0, and one checks that any orbit for $-\mathcal{X}$ can enter neighborhoods $V(K_0)$ of one of these critical submanifolds only a finite number of times, since F is bounded below and decreases by a fixed amount each time. Thus one can show that $\mathcal{M}$ is a disjoint union of the unstable manifolds of the critical points of F.

We remark that the gradient-like vector field $\mathcal{X}$ can be chosen to be tangent to each of the submanifolds $L^2_k(S^2, M)$, for $k \in \mathbb{N}$, $k \geq 2$, because we can choose local pseudogradients and the local representatives $\mathcal{X}_A$ near each critical submanfiold to satisfy this property. Indeed, the only place in the argument where we need the $L^{2\alpha}_1$ topology is for convergence of trajectories to an $L^{2\alpha}_1$ critical point as $t \to \infty$. Then elliptic bootstrapping shows that the critical point is in fact C^∞, and convergence holds in all the Sobolev norms. We formalize this in a remark:

Remark 4.1. We can choose our pseudogradient so that all critical points and all orbits between critical points lie within $L^2_k(S^2, M)$, for any $k \geq 2$.

5 Compactness

We say that a Morse-Bott function $F : \mathcal{M} \to \mathbb{R}$ is of *finite type* if the set of critical points of index λ,

$$K_\lambda = \{f \in \mathcal{M} : dF(f) = 0 \text{ and the Morse index of } f \text{ equals } \lambda \ \}$$

is compact for every $\lambda \in \{0\} \cup \mathbb{N}$. This specializes to the condition that a Morse function have finitely many critical points of each index, or that an equivariant Morse function have only finitely many critical orbits. Thus, for example, if the fundamental group of a compact Riemannian manifold M has infinitely many conjugacy classes, the action integral

$$J : \mathrm{Map}(S^1, M) \longrightarrow \mathbb{R}, \qquad J(\gamma) = \frac{1}{2}\int_{S^1} |\gamma'(t)|^2 dt,$$

is not of finite type, because there are infinitely many local minima, critical points of Morse index zero.

Compactness Theorem 5.1. *a. Suppose that the compact Riemannian manifold M has positive complex sectional curvature, or more generally, positive half isotropic sectional curvature. Then*

$$E_\alpha : L_1^{2\alpha}(S^2, M) \longrightarrow [0, \infty)$$

is of finite type, and indeed there is a constant $c > 0$ such that if $f : S^2 \to M$ is any harmonic or α-harmonic map with $\alpha > 1$, then

$$(\textit{Morse index of } f) + 1 \geq cE(f).$$

b. When M has positive half-isotropic sectional curvature, the set K of critical points for E_α which have Morse index $\leq \lambda_0$ is compact, for any $\lambda_0 \in \mathbb{N}$.

Proof of a when f is harmonic: We start with the case of a harmonic map $f : S^2 \to M$, and note that by a theorem of Koszul and Malgrange, the bundle $\mathbf{E} = f^*TM \otimes \mathbb{C}$ has a canonical holomorphic structure, and the equation for harmonic maps implies that $\partial f/\partial z$ is a holomorphic section of this bundle. Moreover, since the Riemannian metric is invariant under the Levi-Civita connection, it extends to a complex bilinear form

$$\langle \cdot, \cdot \rangle : \mathbf{E} \times \mathbf{E} \longrightarrow \mathbb{C},$$

which is holomorphic as a section of $\mathrm{Hom}(\mathbf{E} \otimes \mathbf{E}, \mathbb{C})$. Let $\mathbf{L}$ be the holomorphic subbundle of $\mathbf{E}$ generated by $\partial f/\partial z$ with orthogonal complement

$$\mathbf{L}^\perp = \bigcup_{p \in S^2} \{v \in \mathbf{E}_p : \langle v, \partial f/\partial z \rangle(p) = 0\}.$$

Then $\mathbf{L} \subseteq \mathbf{L}^\perp$, so we can define the holomorphic normal bundle to be

$$\mathbf{N} = \mathbf{L}^\perp/\mathbf{L}, \tag{27}$$

and $\langle \cdot, \cdot \rangle$ descends to a holomorphic complex bilinear form on $\mathbf{N}$. By a well-known theorem of Grothendieck [14], we can write $\mathbf{N}$ as a holomorphic direct sum of line bundles,

$$\mathbf{N} = \mathbf{L}_1 \oplus \mathbf{L}_2 \oplus \cdots \oplus \mathbf{L}_{n-2}, \text{ where } c_1(\mathbf{L}_1)[S^2] \geq c_1(\mathbf{L}_2)[S^2] \geq \cdots \geq c_1(\mathbf{L}_{n-2})[S^2].$$

Moreover, the existence of a holomorphic complex bilinear form implies that

$$c_1(\mathbf{L}_i)[S^2] = -c_1(\mathbf{L}_{n-i-1})[S^2].$$

The second variation formula (5) for E shows that

$$d^2E(f)(V, \bar{V}) \leq 2 \int_{S^2} \left[\left\| \bar{\partial} V \right\|^2 - \kappa e(f) \langle V, \bar{V} \rangle \right] dA, \quad \text{for} \quad V \in \Gamma(\mathbf{N}),$$

where κ is a positive lower bound on the half-isotropic curvature. Of course, we can restrict to sections of one of the line bundle summands $\mathbf{L}_i$ of $\mathbf{N}$, obtaining

$$d^2E(f)(V,\bar{V}) \leq \int_{S^2} \left[2\left\|\bar{\partial}V\right\|^2 - \phi\langle V,\bar{V}\rangle \right] dA, \quad \text{for} \quad V \in \Gamma(\mathbf{L}_i),$$

where $\phi = 2\kappa e(f) > 0$. To estimate the index of the restriction to $\Gamma(\mathbf{L}_i)$, we need to estimate the spectrum of the Schrödinger operator

$$2\bar{\partial}^* \circ \bar{\partial} - \phi : C^\infty(\mathbf{L}_i) \to C^\infty(\mathbf{L}_i),$$

where $\phi > 0$. An advantage is that the spectrum of this last operator is actually independent of the Hermitian metric:

Lemma 5.2. *A change in the Hermitian metric in the bundle $\mathbf{L}_i$ does not change the spectrum of the operator $2\bar{\partial}^* \circ \bar{\partial} - \phi$.*

To prove this, we need only determine how multiplication of the fiber metric $\langle \cdot, \cdot \rangle$ by a scale factor $\mu^2 > 0$ alters the eigenvalue equation:

$$\int_{S^2} \langle (2\bar{\partial}^* \circ \bar{\partial} - \phi)V, \bar{W}\rangle dA = \lambda \int_{S^2} \langle V, \bar{W}\rangle dA, \quad \text{for } W \in \Gamma(\mathbf{L}_i).$$

But integrands on both sides get multiplied by μ^2 which leads to the eigenvalue equation

$$\int_{S^2} \langle (\bar{\partial}^* \circ \bar{\partial} - \phi)V, \bar{W}\rangle \mu^2 dA = \lambda \int_{S^2} \langle V, \bar{W}\rangle \mu^2 dA.$$

Thus we need find the eigenvalues of the operator

$$T_{\mu^2} \circ \left(2\bar{\partial}^* \circ \bar{\partial} - \phi\right) \circ T_{\mu^2}^{-1},$$

where T_{μ^2} denotes multiplication by the function $\mu^2 > 0$, and conjugation by multiplication by a nonzero function preserves the eigenvalues. QED

The eigenvalues of $2\bar{\partial}^* \circ \bar{\partial} - \phi$ are in turn related to the eigenvalues of a scalar multiple of the operator $2\bar{\partial}^* \circ \bar{\partial}$ by the following reduction, which comes from Li and Yau [21]: Since

$$\frac{\int_{S^2}[2|\bar{\partial}V|^2 - \phi|V|^2]dA}{\int_{S^2}|V|^2 dA} = \frac{\int_{S^2}\phi|V|^2 dA}{\int_{S^2}|V|^2 dA}\left[\frac{\int_{S^2} 2|\bar{\partial}V|^2 dA}{\int_{S^2}\phi|V|^2 dA} - 1\right], \tag{28}$$

the index of the restriction of $d^2E(f)$ to $\Gamma(\mathbf{L}_i)$ is the maximal dimension of a linear subspace of $C^\infty(\mathbf{L}_i)$ on which the quotient

$$V \quad \mapsto \quad \frac{\int_{S^2} 2|\bar{\partial}V|^2 dA}{\int_{S^2}\phi|V|^2 dA}$$

is strictly less than one. We now divide into cases:

$c_1(\mathbf{L}_i) = 0$**:** In this case, we can choose the unitary structure on $\mathbf{L}_i$ so that

$$\frac{\int_{S^2} 2|\bar\partial V|^2 dA}{\int_{S^2} \phi |V|^2 dA} = \frac{\int_{S^2} |dV|^2 dA}{\int_{S^2} \phi |V|^2 dA}.$$

But the right-hand side is just the Rayleigh quotient for the Laplace-Beltrami operator Δ_ϕ for the Riemannian metric

$$\phi ds_0^2, \quad \text{where } ds_0^2 \text{ is the base Riemannian metric on } S^2.$$

It follows from Korevaar's theorem [18] that there is a constant $C > 0$ such that the k-th eigenvalue of Δ_ϕ is less than or equal to one if

$$C\frac{k}{\int_{S^2} \phi dA} \leq 1, \quad \text{which occurs if} \quad k \leq \frac{1}{C}\int_{S^2} \phi dA.$$

Alternatively, we can say that the number of eigenvalues which is less than one is the greatest integer less than

$$C_1 \int_{S^2} \phi dA \quad \text{where} \quad C_1 = \frac{1}{C}.$$

This implies that the index of the Schrödinger operator $2\bar\partial^* \circ \bar\partial - \phi$ grows linearly with the integral of ϕ, which gives the estimate of Theorem 5.1 in this case.

$c_1(\mathbf{L}_i) \neq 0$**:** (Since the sum of the Chern classes is zero, we need only treat the case in which $c_1(\mathbf{L}_i) > 0$.) To treat the case of Chern class $k > 0$, we use the fact that the spectrum of the Laplace operator is not affected much by what happens in the ε-ball around a single point, and note that if we take a single point out of the base of a $U(1)$-bundle, we lose the first Chern class, a fact that we can exploit. Indeed, suppose that $\mathbf{E}_0$ is the trivial line bundle with trivial holomorphic structure. We can let $\mathbf{E} = \mathbf{E}_0 \otimes \nu_p^k$, where ν_p is the point bundle at p (as described for example in [15]), the resulting bundle having a preferred holomorphic section ϕ_0 which has divisor $(\phi_0) = kp$. We can then define a map

$$\iota : C^\infty(S^2, \mathbf{E}_0) \longrightarrow C^\infty(S^2, \mathbf{E}) \quad \text{by} \quad \iota(\phi) = \phi\phi_0. \tag{29}$$

Finally, we let A_0 be a unitary connection on the $U(1)$-bundle $\mathbf{E}$ which has constant curvature and has ϕ_0 as a holomorphic section. We claim that if the m-th eigenvalue λ_m^0 of the operator $2(\bar\partial^* \circ \bar\partial)$ on $\mathbf{E}_0$ is $\leq C$, then so is the m-th eigenvalue λ_m of the operator $2\bar\partial_{A_0}^* \circ \bar\partial_{A_0}$ on $\mathbf{E}$.

Indeed, if $\lambda_m^0 \leq C$, then there is an m-dimensional subspace $\mathcal{V}$ of $C^\infty(S^2, \mathbf{E}_0)$ such that

$$\sigma \in \mathcal{V} \quad \Rightarrow \quad \frac{\int_{S^2} \langle \bar\partial\sigma, \bar\partial\sigma\rangle dA}{\int_{S^2} |\phi|^2 dA} \leq C.$$

We now look at the Rayleigh quotients of the corresponding elements $\iota(\sigma)$ in $C^\infty(S^2, \mathbf{E})$ and exploit the fact that

$$\bar\partial_{A_0}(\sigma\sigma_0) = \bar\partial(\sigma)\sigma_0.$$

This implies that the corresponding Rayleigh quotient in $\mathbf{E}$ is

$$\frac{\int_{S^2}\langle\bar\partial\sigma,\bar\partial\sigma\rangle|\sigma_0|^2dA}{\int_{S^2}|\phi|^2|\sigma_0|^2dA}.$$

But this is just the Rayleigh quotient on the trivial bundle, with the unitary structure multiplied by $|\sigma_0|^2$. This would have the same spectrum as the standard unitary structure, except for the fact that $|\sigma_0|^2$ vanishes at the point p. Indeed, we can approximate $|\sigma_0|^2$ by a nowhere zero function μ^2 in such a way that the support of $|\sigma_0|^2-\mu^2$ is contained in an ε-ball around p. Multiplication by a suitable cutoff function shows that this alters the spectrum by a small amount which goes to zero as $\varepsilon\to 0$, as we next describe.

Construction of a cutoff function: Following Choi and Schoen [8], we define a map $\phi_\varepsilon:\mathbb{R}\to\mathbb{R}$ by

$$\phi_\varepsilon(r)=\begin{cases}0, & \text{if } r\le\epsilon^2,\\ 2-(\log r)/(\log\epsilon), & \text{if } \epsilon^2\le r\le\epsilon,\\ 1, & \text{if } \epsilon\le r,\end{cases}\tag{30}$$

so that

$$\frac{d\phi_\varepsilon}{dr}(r)=\begin{cases}0, & \text{if } r\le\epsilon^2,\\ (-1)/(r\log\epsilon), & \text{if } \epsilon^2\le r\le\epsilon,\\ 0, & \text{if } \epsilon\le r,\end{cases}$$

and

$$\int_0^{2\pi}\int_0^{\epsilon}\left(\frac{d\phi_\varepsilon}{dr}(r)\right)^2 rdrd\theta=\int_{\epsilon^2}^{\epsilon}\frac{2\pi}{r(\log\epsilon)^2}dr=-\frac{2\pi}{\log\epsilon}.$$

Thus if we define $\psi_\varepsilon:\Sigma\to\mathbb{R}$ so that it is one outside an ϵ-neightborhood of the p and in terms of geodesic polar coordinates (r,θ) about p (for the constant curvature metric) satisfies the condition $\psi_\varepsilon=\phi_\varepsilon\circ r$, then

$$\int_{S^2}|d\psi_\varepsilon|^2dA\le\frac{-C}{\log\epsilon},\qquad\text{where } C \text{ is a positive constant,}$$

and ψ_ε is our desired cutoff function which vanishes in the ε^2 neighborhood of the point p, and yet alters Rayleigh quotients by arbitrarily little.

Proof of a when f is α-harmonic: If $f:S^2\to M$ is α-harmonic, the second variation formula (17) for E_α shows that

$$d^2E_\alpha(f)(V,\bar V)\le\alpha\int_{S^2}\left[2\left\|\bar\partial V\right\|^2-\kappa e(f)\langle V,\bar V\rangle\right]dA,\quad\text{for } V\in\Gamma(f^*TM\otimes\mathbb{C}),\tag{31}$$

where κ is a lower bound on the half-isotropic sectional curvature. We now apply the preceding argument to show that the right-hand side of (31) has index which grows linearly with the energy.

Proof of b: if f is a critical point for E_α which has Morse index $\leq \lambda_0$, then it follows from a that $E(f) \leq (\lambda_0 + 1)/c$, or

$$f \in \mathcal{M}^a = \{f \in L_1^{2\alpha}(S^2, M) : E_\alpha(f) \leq a\}, \quad a = \frac{\lambda_0 + 1}{c}.$$

It follows from Condition C that the set of critical points lying within $\mathcal{M}^a$ is compact.

6 Bubble trees and the Nonbubbling Theorem

We are interested in critical points for the usual energy E, not the α-energy E_α or the Morse approximation F_α to E_α defined by (16). Thus we study what happens to the critical points as the perturbations are turned off, for example as $\alpha \to 1$.

This is a special case of a more general question asked by Sacks and Uhlenbeck in [32]. If Σ is a compact Riemann surface with conformal structure ω, the ω-energy is the function

$$E_\omega : \mathrm{Map}(\Sigma, M) \longrightarrow M \quad \text{defined by} \quad E_\omega(f) = \frac{1}{2}\int_\Sigma |df|^2 dA,$$

where $|df|$ and dA are defined in terms of a Riemannian metric in the conformal equivalence class determined by ω, and it has a perturbation

$$E_{\alpha,\omega} : \mathrm{Map}(\Sigma, M) \longrightarrow M \quad \text{defined by} \quad E_{\alpha,\omega}(f) = \frac{1}{2}\int_\Sigma (1 + |df|^2)^\alpha dA - \frac{1}{2},$$

where we lose conformal invariance, and the metric on Σ has constant curvature and total area one.

Suppose that for $m \in \mathbb{N}$, f_m is a critical point for $E_{\alpha_m,\omega}$ with energy bounded by $E_0 > 0$ and Morse index bounded by $\lambda_0 \in \mathbb{N}$. Does the sequence $\{f_m : m \in \mathbb{N}\}$ have a subsequence which has a harmonic map limit as $m \to \infty$? Sacks and Uhlenbeck show that there is such a subsequence which converges in a weak sense to a base harmonic map $f_0 : S^2 \to M$ together with a collection of harmonic two-spheres which bubble off in the limit. Later Parker and Wolfson introduced the term of "bubble tree" to describe the limit, and the nature of the convergence was further studied by Parker [31], and Chen and Tian [6].

Definition. A *bubble tree* based upon a compact Riemann surface Σ_g of genus $g \geq 0$ with conformal structure ω is a collection of maps indexed by the vertices and edges of a rooted tree T, with root denoted by 0. The root of the tree corresponds to an ω-harmonic *base map*

$$f_0 : \Sigma_g \to M, \quad \text{when } v = 0,$$

and the other vertices correspond to harmonic two-spheres or *bubble maps*

$$f_v : S^2 \to M, \quad \text{when } v \neq 0,$$

while the edges correspond to smooth geodesics called *necks*

$$\gamma_v : [0,1] \to M$$

which join the parent $p(v)$ of a vertex to a vertex v (with γ_v only defined when $v \neq 0$), so that

$$\gamma_v(0) \in \text{Image}(f_{p(v)}), \quad \gamma_v(1) \in \text{Image}(f_v).$$

We allow a vertex v to correspond to a constant map, but only when v is the parent of another vertex. In this case we say that f_v is a *ghost bubble*. We also allow some of the necks γ_v to be constant maps to a single point of M.

If we assume an upper bound on total energy, we get a limit on the number of possible bubbles from the following elementary lemma:

Lemma 6.1. *Suppose that the real sectional curvatures of M satisfy the inequality $K_r(\sigma) \leq 1$. Then any nonconstant minimal two-sphere $f : S^2 \to M$ must have area at least 4π.*

Proof: It follows from the Gauss equation,

$$\text{induced Gaussian curvature } K_f \text{ on } S^2 \leq K_r((f_p)_* T_p S^2)$$

and the Gauss-Bonnet formula that if $f : S^2 \to M$ is any minimal two-sphere,

$$E(f) \geq \int K_f dA_f \geq 4\pi,$$

the last inequality being strict if f has nontrivial branch locus.
Hence if the total energy is bounded above by $a > 0$, the number of minimal two-spheres that can form as bubbles is $\leq a/4\pi$.

Construction of a bubble tree: Suppose that $\{f_m\}$ is a sequence of critical points for $E_{\alpha_m,\omega}$ with $m \to \infty$ such that $\alpha_m \to 1$, $E_{\alpha_m,\omega}(f_m)$ is bounded by some constant $a > 0$, and $f_m : \Sigma \to M$ has bounded Morse index. According to Theorem 4.4 of Sacks and Uhlenbeck [32], $\{f_m\}$ has a subsequence (denoted again by $\{f_m\}$ for simplicity) which converges in C^k for all k on compact subsets of the complement in Σ of a finite subset $\{p_1, \ldots, p_l\}$ of "bubble points" to a smooth ω-harmonic map

$$f_0 : \Sigma \to M.$$

For each m, we divide the Riemann surface Σ into a collection of metric disks $D_{1;m}, \ldots, D_{l;m}$ of small radius about the bubble points $\{p_1, \ldots, p_l\}$ and a *base*

$$\Sigma_{0;m} = \Sigma - (D_{1;m} \cup \cdots \cup D_{l;m}).$$

Each disk $D_{i;m}$ is further decomposed into a union $D_{i;m} = A'_{i;m} \cup B'_{i;m}$, where $A'_{i;m}$ is an annular neck region and $B'_{i;m}$ is a bubble region, a smaller disk which

is centered at p_i. Following Parker [31], we arrange that the radius of $D_{i;m}$ is $\leq$ (constant)$/m$, the radius of $B'_{i;m}$ is $\leq$ (constant)$/m^3$, and

$$\frac{1}{2}\int_{A'_{i;m}} |df_m|^2 dA = C_R,$$

where C_R is a small positive renormalization constant, too small to allow bubbling in the annulus $A'_{i;m}$. Let $B_{i;m}$ be a disk with the same center as $B'_{i;m}$ but with m times the radius and let $A_{i;m} = D_{i;m} - B_{i;m}$. If the renormalization constant is sufficiently small, the supremum of $|df_m|$ on $A_{i;m}$ must go to zero because of the Sacks-Uhlenbeck ϵ-regularity theorem (see [32], Main Estimate 3.2 or [31]).

We expand the disk $B_{i;m}$ to a disk of radius m by means of a conformal contraction $T_{i;m} : D_m(0) \to B_{i;m}$, $D_m(0)$ being the disk of radius m in $\mathbb{C}$. An argument based upon Theorem 4.4 of [32] shows that a subsequence of

$$g_{i;m} = f_m \circ T_{i;m} : D_m(0) \to M, \qquad m = 1, 2, \ldots$$

converges uniformly in C^k on compact subsets of $\mathbb{C}$, or on compact subsets of $\mathbb{C} - \{p_{i,1}, \ldots, p_{i,l_i}\}$, where $\{p_{i,1}, \ldots, p_{i,l_i}\}$ is a finite set of new bubble points, to a harmonic map $g_i : S^2 \to M$.

If there are new bubble points $\{p_{i,1}, \ldots, p_{i,l_i}\}$ in $\mathbb{C} = S^2 - \{\infty\}$, the process can be repeated. Around each new bubble point $p_{i,j}$, we construct a small disk $D_{i,j;m}$ which is further subdivided into an annular region $A_{i,j;m}$ and a smaller disk $B_{i,j;m}$ on which bubbling will occur. Once again, we construct conformal contractions $T_{i,j;m} : D_m(0) \to B_{i,j;m}$ and a subsequence of the maps $g_{i,j;m} = f_m \circ T_{i,j;m}$ will converge to a harmonic two-sphere $g_{i,j} : S^2 \to M$ on the complement of finitely many points. The process can be repeated; for each f_m in the sequence, we may have several *level-one* bubble regions $B_{i;m}$, each of which may contain several *level-two* bubble regions $B_{i,j;m}$, each of which may contain several *level-three* bubble regions, and so forth. For each sufficiently large $m \in \mathbb{N}$, Σ is a disjoint union of a *base* $\Sigma_{0;m}$, *neck* regions $A_{i_1,\ldots,i_k;m}$ and *bubble* regions

$$\Delta_{i_1,\ldots,i_k;m} = B_{i_1,\ldots,i_k;m} - \bigcup_j D_{i_1,\ldots,i_k,j;m}, \tag{32}$$

from which disks around bubble points of higher level have been deleted. The restriction $g_{i_1,\ldots,i_k;m}$ of f_m to the bubble regions $\Delta_{i_1,\ldots,i_k;m}$ corresponding to a given multiindex $(i_1, \ldots i_k)$ converge to a harmonic two-sphere $g_{i_1,\ldots,i_k}$ which may have zero energy (in which case it is called a *ghost bubble*), but ghost bubbles always have bubble points in addition to ∞. Since each bubble with no children is nonconstant and requires energy $\geq 4\pi$, the process must terminate after finitely many steps, yielding harmonic maps corresponding to the vertices of a finite tree T (a base ω-harmonic surface $g_0 : \Sigma \to M$ and several minimal two-spheres), while the edges can be thought of as limits of parametrized maps from the annular regions into M.

Lower Semicontinuity Theorem 6.2. *The Morse indices of the components of the bubble tree satisfy:*

$$\text{Morse index of } f_0 + \sum(\text{Morse index of } g_{i_1,\dots i_k}) \le \text{Morse index of } f_m, \quad (33)$$

when m is sufficiently large.

Proof: We first recall the second variation formula for α-harmonic maps f:

$$d^2E_{\alpha,\omega}(f)(V,W) = \alpha\int_\Sigma (1+|df|^2)^{\alpha-1}[\langle\nabla V,\nabla W\rangle - \langle\mathcal{K}(V),W\rangle]dA \\ + 2\alpha(\alpha-1)\int_\Sigma (1+|df|^2)^{\alpha-2}\langle df,\nabla V\rangle\langle df,\nabla W\rangle dA.$$

Note that the second term is dominated by the first and goes to zero as $\alpha\to 1$, and the first term approaches $d^2E_\omega(f)(V,W)$ when the supports of V and W do not contain any bubble points.

We claim that if $d^2E_\omega(f_0)$ or $d^2E(g_{i_1,\dots i_k})$ is negative definite on a linear space $\mathcal{V}$ of dimension m, so is

$$d^2E_{\alpha_m,\omega}(f_m|\Sigma_{0;m}) \quad \text{or} \quad d^2E_{\alpha_m,\omega}(f_m|\Delta_{i_1,\dots,i_k;m})$$

when m is sufficiently large. Since $\Sigma_{0;m}$ and the bubble regions $\Delta_{i_1,\dots,i_k;m}$ are all mutually disjoint, this will prove (33).

To establish our claim in the case of f_0, we make use of a cutoff function defined utilizing the function ϕ_ε of (30) near the bubble points. We define $\psi_i : S^2\to\mathbb{R}$ so that it is one outside an ϵ-neightborhood of the bubble point x_i and in terms of polar coordinates (r_i,θ_i) about the bubble point x_i satisfies the condition $\psi_i = \phi_\varepsilon\circ r_i$, so that

$$\int_{S^2}|d\psi_i|^2dA \le \frac{-C}{\log\epsilon}, \qquad \text{where } C \text{ is a positive constant,}$$

and check that a similar estimate holds for $\psi=\psi_1\cdots\psi_l$, a cutoff function which vanishes at every bubble point. If $\varepsilon>0$ is chosen sufficiently small, then

$$V\in\mathcal{V} \quad\Rightarrow\quad d^2E(f)(\psi V,\psi V)<0 \quad\Rightarrow\quad d^2E_\alpha(f)(\psi V,\psi V)<0.$$

This shows that $d^2E_\alpha(f)$ is negative definite on a space of dimension m, establishing the claim in this case.

Similarly, we can establish the claim in the case of one of the minimal two-spheres $g_{i_1,\dots i_k}$. In this case, the rescalings of the bubble regions $\Delta_{i_1,\dots,i_k;m}$ have boundary circles which converge to points, which we call the bubble points of $g_{i_1,\dots i_k}$, and $g_{i_1,\dots i_k}$ is defined on the sphere S^2 minus small disks around bubble points which converge to punctured disks in S^2. But the cutoff argument in the preceding paragraph can still be applied and this proves the Lemma.

The Nonbubbling Theorem 1.4 follows from Theorem 6.2 and the fact that any harmonic two-sphere in a manifold with positive complex sectional curvatures

has Morse index at least $n-2$. Thus if a bubble tree contains k bubbles of nonzero area, it must have Morse index at least $k(n-2)$.

Compactness Theorem 6.3. *Suppose that M has positive half-isotropic sectional curvatures and all nonconstant minimal two-spheres $f : S^2 \to M$ of Morse index $\leq 2n-5$ lie on Morse nondegenerate $\mathcal{G}$-orbits, where $\mathcal{G} = PSL(2,\mathbb{C}) \cup R \cdot PSL(2,\mathbb{C})$. Then there are only finitely many such orbits of Morse index $\leq 2n-5$.*

Proof: Suppose the contrary, that there is a sequence $\{f_m\}$ of minimal two-spheres lying on distinct G-orbits. Then we can use the Perturbation Theorem 2.2 to construct a corresponding sequence $\{f'_m\}$ of α_m-energy critical points for E_{α_m}, with $\alpha_m \to 1$. By the Compactness Theorem 5.1, these must all have energy $\leq c$, where c depends on the maximum possible Morse index $2n-5$. Now a subsequence of these must converge to a harmonic map $f_\infty : S^2 \to M$ without bubbling. The reason no bubbling can occur is that this would contradict the Nonbubbling Theorem 1.1. But then we have a sequence of harmonic maps on distinct G-orbits which converges to f_∞ contradicting the nondegeneracy of the limit.

7 Branched covers

Suppose that $h : S^2 \to M$ is an immersed prime minimal two-sphere. If $g : S^2 \to S^2$ is a holomorphic map (also known as a meromorphic function on S^2), then the composition $f = h \circ g : S^2 \to M$ is a conformal harmonic map, hence a minimal surface in its own right, which we call a *branched cover* of h. To study these, we consider the space

$$\mathrm{Hol}_d(S^2, S^2) = \{g : S^2 \to S^2 : \ g \text{ is holomorphic and has degree } d \ \},$$

for each integer $d \geq 2$. The topology of the spaces $\mathrm{Hol}_d(S^2, S^2)$ of holomorphic maps, the more general spaces spaces $\mathrm{Hol}_d(\Sigma, S^2)$ of meromorphic functions on a Riemann surface Σ of higher genus, and how well these spaces approximate the spaces of all maps of degree d, were studied by Segal [34].

We can construct elements of $\mathrm{Hol}_d(S^2, S^2)$ as follows: Take two effective divisors D_0 and D_∞ on S^2 of degree d, which have no points in common. Then there will be a unique meromorphic function g on S^2 with divisor $(g) = D_0 - D_\infty$, up to multiplication by a nonzero complex scalar. In fact, if we regard S^2 as $\mathbb{C} \cup \{\infty\}$, and arrange after a conformal automorphism of S^2 that neither divisor includes infinity, say

$$D_0 = \sum_{i=1}^{d} p_i, \quad D_\infty = \sum_{i=1}^{d} q_i, \quad \text{then} \quad g = c\frac{(z-p_1)\cdots(z-p_d)}{(z-q_1)\cdots(z-q_d)}$$

is a corresponding meromorphic function, where the complex number c can be an arbitrary element of $\mathbb{C} - \{0\}$. (Note that we allow repetition among the p_i's

and q_i's.) From this description, we see that $\mathrm{Hol}_d(S^2, S^2)$ is a complex manifold of complex dimension $2d+1$,

$$\mathrm{Hol}_d(S^2,S^2) = (\mathbb{C}-\{0\}) \times \left[\overbrace{S^2 \times \cdots \times S^2}^{d} \times \overbrace{S^2 \times \cdots \times S^2}^{d} - \hat{\Delta} \right],$$

where $\hat{\Delta}$ is the multidiagonal,

$$\hat{\Delta} = \{(p_1, \ldots, p_d, q_1, \ldots, q_d) : \text{some } p_i \text{ equals some } q_j\}.$$

We see that $\mathrm{Hol}_d(S^2, S^2)$ is noncompact, and in fact if one of the p_i's moves into coincidence with one of the q_j's, one can check that the L^2_k norm of g must go to infinity, when $k \geq 2$.

For each integer $d \geq 1$, we define a map

$$\Phi_d : \mathrm{Map}(S^2, M) \times \mathrm{Hol}_d(S^2, S^2) \to \mathrm{Map}(S^2, M) \quad \text{by} \quad \Phi_d(h, g) = h \circ g,$$

a map which will be C^∞ when the mapping spaces are completed with respect to the L^2_k topology, according to the α-lemma on L^2_k spaces. If h is an immersed prime minimal surface, the image of Φ_d is a critical submanifold

$$\mathcal{M}_d(h) \subseteq L^2_k(S^2, M).$$

However, if $\mathcal{N}$ is a finite-dimensional critical submanifold of $L^2_k(S^2, M)$, nondegenerate in the sense of Bott, examples show that the union of d-fold branched covers of $\mathcal{N}$,

$$\Phi_d(\mathcal{N} \times \mathrm{Hol}_d(S^2, S^2)) = \bigcup \{\mathcal{M}_d(h) : h \in \mathcal{N}\},$$

may contain critical points of various indices and hence is not necessarily a nondegenerate critical submanifold in the sense of Bott.

We can ask how the Morse index of the branched cover varies over this submanifold, and obtain the following estimate:

Theorem 7.1. *Suppose M is an n-dimensional Riemannian manifold which has isotropic and real sectional curvatures which satisfy the curvature condition*

$$K_i(\sigma) > \frac{1}{d} K_r(\widehat{\sigma}) > 0, \tag{34}$$

where $\widehat{\sigma}$ is a real two-plane associated to σ. Then the Morse index of any branched cover of degree $\geq d$ of a minimal two-sphere $f : S^2 \to M$ must be at least $2(n-2)$.

Note that the curvature condition reduces to (7) when $d = 3$.

Proof of theorem: As in the proof of Theorem 5.1, we use the normal bundle (27) and its holomorphic decomposition into line bundles,

$$\mathbf{N} = \mathbf{L}_1 \oplus \mathbf{L}_2 \oplus \cdots \oplus \mathbf{L}_{n-2}, \text{ where } c_1(\mathbf{L}_1)[S^2] \geq c_1(\mathbf{L}_2)[S^2] \geq \cdots \geq c_1(\mathbf{L}_{n-2})[S^2].$$

If $V = X - iY$ is a holomorphic section of a line bundle summand $\mathbf{L}_i$ with $c_1(\mathbf{L}_i)[S^2] > 0$, then V must vanish at some point, and since $\langle V, V\rangle$ is a bounded holomorphic function and hence constant, $\langle V, V\rangle \equiv 0$. This implies that the real and imaginary components X and $-Y$ must have the same length and be perpendicular to each other. If E_i denotes the rank two real subbundle of f^*TM generated by real and imaginary parts of sections of $\mathbf{L}_i$, we can define an almost complex structure J on E_i by

$$J(X) = Y, \quad \text{whenever} \quad V = X - iY \in \Gamma(\mathbf{L}_i).$$

Then the components X and Y of a holomorphic section $V = X - iY$ of $\mathbf{L}_i$ generate a two-dimensional subspace of sections of E_i, complex linear with respect to J, on which the Morse index is negative. Since

$$c_1(\mathbf{L}_i)[S^2] = k > 0 \quad \Rightarrow \quad \dim \mathcal{O}(\mathbf{L}_i) = k + 1,$$

the corresponding real and imaginary parts form a linear subspace of sections of real dimension $2k + 2$ on which the index form is negative. The upshot is that each summand with positive Chern class must contribute at least four to the Morse index.

If there are m summands with positive first Chern class (where $m \leq (n - 2)/2$), there will be $n - 2 - 2m$ summands with zero Chern class. Thus if we can show that each zero Chern class summand contributes at least two to the Morse index, the total Morse index will be at least

$$4m + 2 \times (n - 2 - 2m) = 2n - 4,$$

establishing Theorem 7.1.

Thus we need only show that our curvature hypothesis (7) implies that trivial line bundle summands must each contribute at least two to the Morse index. For this purpose, we can apply the second variation formula (5) which shows that

$$\begin{aligned} d^2E(f)(V, \bar{V}) &= 2\int_{S^2} \left[\left\| \bar{\partial} V \right\|^2 - K_i(\sigma) e(f) \left\| V \right\|^2 \right] dA \\ &= \int_{S^2} \left[2 \left\| \bar{\partial} V \right\|^2 - 2K_i(\sigma) \left\| V \right\|^2 \right] d\tilde{A}, \quad (35) \end{aligned}$$

where $d\tilde{A} = e(f)dA$ is the area form for the induced metric on S^2, and σ is the span of V and $\frac{\partial f}{\partial z}$. From (35), we conclude that

$$d^2E(f)(V, \bar{V}) = \int_{S^2} \left[\left\| dV \right\|^2 - 2K_i(\sigma) \left\| V \right\|^2 \right] d\tilde{A},$$

because by Lemma 5.2, we can assume that the connection is flat. Substituting

the curvature condition (34), we have

$$d^2E(f)(V,\bar{V}) < \int_{S^2}\left[\|dV\|^2 - \frac{2}{d}K_r(\widehat{\sigma})\,\|V\|^2\right]d\tilde{A},$$
$$= \left(\frac{\int_{S^2}\|dV\|^2\,d\tilde{A}}{\int_{S^2}K_r(\widehat{\sigma})\,\|V\|^2\,d\tilde{A}} - \frac{2}{d}\right)\int_{S^2}K_r(\widehat{\sigma})\,\|V\|^2\,d\tilde{A},$$
$$\leq \left(\lambda_1 - \frac{2}{d}\right)\int_{S^2}K_r(\widehat{\sigma})\,\|V\|^2\,d\tilde{A}, \quad (36)$$

where λ_1 is the first nonzero eigenvalue of the Laplace-Beltrami operator for the metric $K_r(\widehat{\sigma})f^*g$ on S^2. Theorem 1 of Yang and Yau [42] implies that λ_1 satisfies

$$\lambda_1 \leq \frac{8\pi}{\int_{S^2}K_r(\widehat{\sigma})d\tilde{A}} \leq \frac{8\pi}{\int_{S^2}K_f d\tilde{A}} \leq \frac{8\pi}{4\pi d} = \frac{2}{d},$$

for a d-fold branched cover, where we have used that $\widehat{\sigma} = f_*TS^2$ and, as in Lemma 6.1, K_f is the Gauss curvature of the induced metric which satisfies, $K_f \leq K_r(f_*TS^2)$. This shows that there are at least two variation fields for each trivial summand of the normal bundle which decrease area, finishing the proof.

One can also use the argument for Theorem 7.1 to establish:

Proposition 7.2. *If $M = S^n$ has the metric of constant curvature one and $h : S^2 \to S^n$ is a totally geodesic imbedding, then the first nonzero eigenvalue for any double branched cover $f \in \mathcal{M}_2(h)$ is ≤ 1.*

The analysis of two-fold branched covers simplifies because any element of $g \in \mathrm{Hol}_2(S^2, S^2)$ can be related to the standard two-fold cover,

$$g = S^{-1} \circ \pi_2 \circ T, \quad \text{where} \quad S, T \in PSL(2,\mathbb{C}) \quad \text{and} \quad \pi_2(z) = z^2. \quad (37)$$

Recalling the discussion on center of mass from §2, we reduce the structure group on $\mathcal{M}_2(h)$ from $\widehat{\mathcal{G}}$ to $\widehat{G} = O(3)$ by restricting to the submanifold

$$\mathcal{M}_2(h)_0 = \left\{ f \in \mathcal{M}_2(h) : \int_{S^2}\mathbf{X}\psi_1(|df|^2)dA = \mathbf{0}\right\}$$

of branched covers with center of mass zero, the action of $\widehat{G} = O(3)$ on $\mathcal{M}_2(h)_0$ preserving the center of mass zero condition. It follows from the explicit form (37) of double branched covers that any element of $\mathcal{M}_2(h)_0$ must have nontrivial isotropy for the action of $G = SO(3)$, the isotropy group being $\mathbb{Z}_2$. Milnor [25] describes the quotient of $\mathcal{M}_2(h)_0$ by the $SO(3)$ action as a four-manifold which is the total space of a two-plane bundle over $\mathbb{R}P^2$.

Although $\mathcal{M}_2(h)_0$ is not compact, we can make it compact by removing the branched covers whose two branch points are too close. Given $\varepsilon_0 > 0$, we set

$$\mathcal{M}_2(h)_0^{\varepsilon_0} = \{ f \in \mathcal{M}_2(h)_0 : \text{ the two branch points have distance} \geq \varepsilon_0 \},$$

the distance being measured in the standard metric of constant curvature and total area one on S^2. We can describe the topological type of this space as follows: For the given fixed prime minimal immersion $h : S^2 \to M$ we can specify a $\widehat{G}$-orbit of branched covers of h by locating the images of its two branch points. Let $n \in S^2$ be the first image, and regard

$$S^2 - (\ \varepsilon_0\text{-neighborhood of } n\ )$$

as a two-disk D_n. The second image is then a point in D_n and the image of the branch locus is specified by a point in a D^2-bundle over S^2. But we can interchange the two branch points and a quotient by the resulting $\mathbb{Z}_2$ action gives us the two-disk bundle N^4 over $\mathbb{R}P^2$. Then $\mathcal{M}_2(h)_0^{\varepsilon_0}$ itself is the total space of a fibration

$$\pi : \mathcal{M}_2(h)_0^{\varepsilon_0} \longrightarrow N^4 \tag{38}$$

the fiber being a quotient space $O(3)/\mathbb{Z}_2$, where $\mathbb{Z}_2$ is a subgroup but not normal. Each element of $\mathcal{M}_2(h)_0^{\varepsilon_0}$ has $\mathbb{Z}_2$ isotropy under the $SO(3)$ action, while the larger $O(3)$ action interchanges the two components of $\mathcal{M}_2(h)_0^{\varepsilon_0}$.

We next study the α-energy critical points which are close to $\mathcal{M}_2(h)_0$, for a given choice of $h : S^2 \to M$. We can use the L^2_k topology where k is large, so our function spaces are Hilbert manifolds, as described in Lang [19]. The Hilbert space inner product on $L^2_k(S^2, \mathbb{R}^N)$ makes it into a flat infinite-dimensional Riemannian manifold as described in Chapter VII of [19], and the isometric imbedding $M \subseteq \mathbb{R}^N$ induces an imbedding $L^2_k(S^2, M) \subseteq L^2_k(S^2, \mathbb{R}^N)$, which induces a Riemannian metric on $L^2_k(S^2, M)$ for which $\widehat{G}$ acts as isometries; this in turn induces a Riemannian metric on the finite-dimensional manifold $\mathcal{M}_2(h)_0$, a space on which $\widehat{G}$ acts with $\mathbb{Z}_2$ isotropy. We now apply the tubular neighborhood theorem: for a suitable $\widehat{G}$-invariant open neighborhood $\nu\mathcal{M}_2(h)_0$ of the zero section in the normal bundle to $\mathcal{M}_2(h)_0$ within $L^2_k(S^2, M)$, we can define a smooth $\widehat{G}$-equivariant diffeomorphism

$$\phi : \nu\mathcal{M}_2(h)_0 \longrightarrow U,$$

where U is some $\widehat{G}$-invariant open tubular neighborhood of $\mathcal{M}_2(h)_0$ in $L^2_k(S^2, M)$. The function E_α restricts to a $\widehat{G}$-invariant function on U.

If $f_0 \in \mathcal{M}_2(h)_0^{\varepsilon_0}$, we let

$$N(f_0) = \text{fiber of } \nu\mathcal{M}_2(h)_0^{\varepsilon_0} \text{ over } f_0,$$

a submanifold of finite codimension within $L^2_k(S^2, M)$, and we restrict the function E_α to this submanifold, $E_\alpha : N(f_0) \to \mathbb{R}$. By an immediate extension of the argument for Theorem 2.2 we can prove:

Proposition 7.3. *There exists $\alpha_0 \in (1, \infty)$ and a smooth map*

$$\eta : \mathcal{M}_2(h)_0^{\varepsilon_0} \times [1, \alpha_0) \longrightarrow \left\{ (f, \alpha) \in L^2_k(S^2, M) \times [1, \alpha_0) : \int_{S^2} \mathbf{X}\psi_\alpha(|df|^2) dA = \mathbf{0} \right\} \tag{39}$$

which satisfies the following conditions:

1. *η is $O(3)$-equivariant,*

2. *$\eta_\alpha(f) = \eta(f,\alpha)$ is the unique critical point for the restriction of E_α to $N(f_0)$ for $f_0 \in \mathcal{M}_2(h)_0^{\varepsilon_0}$ and $\alpha \in [1,\alpha_0)$, and this critical point is nondegenerate on $N(f_0)$,*

3. *there is an $\varepsilon_1 > 0$ such that if f is a critical point for E_α such that $\|f - f_0\|_{C^2} < \varepsilon_1$ for some $f_0 \in \mathcal{M}_2(h)_0^{\varepsilon_0+\varepsilon_1}$, f lies in the image of η, and*

4. *the isotropy group under the $SO(3)$ action of every element in the image of η is $\mathbb{Z}_2$, a rotation through angle π about some axis.*

For the third condition on η, note that if f is an element of $N(f_0)$ and a critical point for E_α, it must also be critical point for the restriction of E_α to $N(f_0)$. This proposition shows that critical points of E_α which are sufficiently close to the submanifold of two-fold branched covers must necessarily be invariant under a rotation in $O(3)$ of 180 degrees.

But we would like those critical points to lie on nondegenerate critical submanifolds, each an orbit for the $\widehat{G}$-action. We can arrange this by first constructing a suitable smooth function $h : N^4 \to \mathbb{R}$, then letting $\tilde{h} = h \circ \pi$, where π is the projection of (38). Since $\tilde{h}$ is constant on each fiber of π it is invariant under the action of $\widehat{G}$. Finally, we let $\psi_\alpha : U \to [0,1]$ be a smooth $\widehat{G}$-invariant function which is identically one on the image of the function $\eta_\alpha = \eta(\cdot,\alpha)$ defined by (39), and let

$$F_h = E_\alpha + \psi_\alpha \tilde{h}, \tag{40}$$

For small h this is a $\widehat{G}$-equivariant perturbation of E_α every critical point of which has $\mathbb{Z}_2$ isotropy under the $\widehat{G}$ action.

We claim that for generic choice of h, the perturbed function F_h will have all critical points lying on isolated $\widehat{G}$ orbits, each nondegenerate in the sense of Bott. Indeed, the only critical points for F_h which lie in $\phi(N(f_0))$ for some $f_0 \in \mathcal{M}_2(h)_0^{\varepsilon_0}$ must be critical points for the restriction of E_α to $\phi(N(f_0))$ and must therefore lie in the image of

$$\eta_\alpha : \mathcal{M}_2(h)_0^{\varepsilon_0} \longrightarrow U.$$

On this image $\psi_\alpha \equiv 1$, and

$$F_h \equiv E_\alpha + \tilde{h},$$

is a small perturbation of E_α when h is small. At a critical point $\eta_\alpha(f_0)$ for F_h, the tangent space divides into a direct sum

$$T_{\eta_\alpha(f_0)} L^2_k(S^2, M) \cong V \oplus H \oplus N, \quad \text{where} \quad H \oplus N \cong T_{f_0}\mathcal{M}_2(h)_0,$$

and we arrange that V is tangent to the fiber of $\nu\mathcal{M}_2(h)_0$ (the "vertical space"), H projects via (38) to the tangent space to N^4 (the "horizontal space"), and that N is tangent to the orbit of the $\widehat{G}$ action. At a critical point, $DE_\alpha = 0$

and $d\tilde{h} = 0$. Invariance under $\widehat{G}$ implies that N is a null-space for both D^2E_α and $d^2\tilde{h}$. Moreover, since h is constant on each fiber of the normal bundle,

$$\nu\mathcal{M}_2(h)_0 \longrightarrow \mathcal{M}_2(h)_0,$$

the space V must lie in the null space of $d^2\tilde{h}$. By part 2 of Proposition 7.3, D^2E_α is nondegenerate on $V \times V$, and a generic choice of $\tilde{h}$ will make

$$d^2F_h = D^2E_\alpha + d^2\tilde{h} \quad \text{nondegenerate on} \quad (V \oplus H) \times (V \oplus H),$$

establishing the claim. We can summarize what we have proven as follows.

Proposition 7.4. *Suppose that $h : S^2 \to M$ is an imbedded minimal two-sphere in the compact Riemannian manifold M and that U is a small open neighborhood of the space $\mathcal{M}_2(h)_0^{\varepsilon_0+\varepsilon_1}$ of two-fold branched covers of h, where ε_0 and ε_1 can be taken arbitrarily small. Then there is a perturbation $\tilde{E}_\alpha$ of E_α which agrees with E_α outside U, is C^2 close to E_α, and satisfies the following condition: All critical points for $\tilde{E}_\alpha$ lie on nondegenerate critical submanifolds, each an orbit for the $\widehat{G}$ action. Moreover, each critical point has isotropy group $\mathbb{Z}_2$, generated by some rotation through π.*

We have left out the points in $\mathcal{M}_2(h)_0 - \mathcal{M}_2(h)_0^{\varepsilon_0}$, but when the half-isotropic curvature is positive, these must have Morse index $\geq 2n-4$ when ε_0 is sufficiently small, by the proof of the Nonbubbling Theorem 1.4.

8 The topology of the mapping space

The strategy for proving the Main Theorem 1.2 is to first use equivariant Morse theory to calculate the low degree equivariant cohomology of the mapping space $\mathrm{Map}(S^2, M)$ (carried out in this section), and then use equivariant Morse theory in the other direction (carried out in § 10) to prove existence of minimal two-spheres in manifolds which satisfy our curvature hypotheses.

Since the α-energy $E_\alpha : \mathrm{Map}(S^2, M) \to \mathbb{R}$ is invariant under an action of the group $O(3)$ by isometries on the domain, the nonconstant critical points for E_α lie on $O(3)$-orbits of critical points, which can be measured by $O(3)$-equivariant Morse theory. We review the essential features of equivariant cohomology needed for this theory.

Any compact Lie group G possesses a classifying space BG together with a universal principal G-bundle with contractible total space EG and right action $\alpha_E : EG \times G \to EG$. If X is a CW complex and $\alpha_X : X \times G \to X$ is a continuous right action, the diagonal action on the product $X \times EG$,

$$(\sigma, (x, e)) \mapsto (\alpha_X(x, \sigma), \alpha_E(e, \sigma)),$$

is free, and we can therefore divide by this action to obtain the topological space X_G; to be more precise we set

$$X_G = (X \times EG)/G,$$

the set of equivalence classes in $X \times EG$ under the equivalence relation

$$(\alpha_X(x,\sigma), \alpha_E(e,\sigma)) \sim (x,e), \qquad \text{for } x \in X,\ e \in EG \text{ and } \sigma \in G.$$

We then define the *equivariant cohomology* of X with coefficients in the coefficient ring R by

$$H^*_G(X;R) = H^*(X_G;R).$$

In particular if X is a point,

$$H^*_G(X;R) = H^*(BG;R).$$

The projection

$$\pi : X_G = X \times_G EG \longrightarrow EG/G = BG$$

defines an action of $H^*(BG;R)$ on $H^*_G(X;R) = H^*(X_G;R)$ making $H^*(X_G;R)$ into a graded module over the graded algebra $H^*(BG;R)$, via

$$(a,b) \in H^*(X_G;R) \times H^*(BG;R) \mapsto a \cup \pi^*(b) \in H^*(X_G;R).$$

This module structure helps measure the isotropy groups. We will apply equivariant cohomology in the case where $G = O(3)$ and the coefficient group is $\mathbb{Z}_2$, and in this case,

$$H^*(BO(3);\mathbb{Z}_2) = P[w_1, w_2, w_3]$$

a polynomial algebra on generators w_1, w_2 and w_3 of dimensions one, two and three, which we can identify with the Stiefel-Whitney classes of the universal bundle over $BO(3)$.

If M is a compact Riemannian manifold, we let M_0 denote the submanifold of $\mathcal{M} = \mathrm{Map}(S^2, M)$ consisting of constant maps, so

$$M_0 = \{f \in L^{2\alpha}_1(S^2, M) : E_\alpha(f) = 0\}$$

is a submanifold consisting entirely of critical points, which is diffeomorphic to M. Let

$$\mathcal{M}^a = \{f \in \mathcal{M} : E_\alpha(f) \le a\}.$$

We can use equivariant Morse theory to describe how the relative cohomology

$$H^*_{O(3)}(\mathcal{M}^a, M_0, \mathbb{Z}_2)$$

changes as a increases. We start the process with:

Lemma 8.1. *There is an $\varepsilon > 0$ such that M_0 is a strong deformation retract of $\mathcal{M}^\varepsilon$.*

This is proven in [32]; see Theorem 2.6. In fact, it is not hard to show that M_0 is a nondegenerate critical submanifold.

Suppose now that E_α is a Morse-Bott function; that is all of its critical points of positive energy lie on compact nondegenerate critical submanifolds. It follows

quickly from Condition C that only finitely many such submanifolds lie within $\mathcal{M}^a$ for $a < \infty$.

We can next establish Theorems A and B as in Bott's presentation [4].

Theorem A. *If the interval $[a, b]$ contains no critical values for E_α, then $\mathcal{M}^a$ is a strong deformation retract of $\mathcal{M}^b$.*

Theorem A is proven by an argument virtually identical to the finite-dimensional case treated in [23]; the deformation retraction is constructed by following the flow lines of $-\mathcal{X}$, where $\mathcal{X}$ is the gradient-like vector field constructed before. We say that c is a critical value for F is there exists a critical point f for F with $F(f) = c$; the following theorem is proven in [37], for a Morse perturbation F of E_α:

Theorem B. *If $[a, b]$ contains a single critical value c and there is exactly one Morse nondegenerate critical point p such that $F(p) = c$, then $\mathcal{M}^b$ is homotopy equivalent to $\mathcal{M}^a$ with a handle of index λ attached.*

But we need a version of Theorem B which applies to E_α, to the case in which we have nondegenerate critical submanifolds instead of nondegenerate critical points. For a nondegenerate critical submanifold $K \subset \mathcal{M}$, the normal bundle to K is defined via the inner product $\langle\langle \cdot, \cdot \rangle\rangle$ defined by (19): If $f \in K$,

$$\nu K_f = \{V \in T_f\mathcal{M} : \langle\langle V, W \rangle\rangle = 0 \text{ for all } W \in T_f K\}.$$

Referring back to the operator A defined before, we see that the normal bundle has a finite-dimensional subbundle $\nu_- K$ generated by eigenvectors corresponding to the negative eigenvalues of A. The rank of the bundle $\nu_- N$ is the *Morse index* of the nondegenerate critical submanifold K. Let

$$D(\nu_- K) = \{V \in \nu_- K : \|V\| \leq 1\}, \qquad S(\nu_- K) = \{V \in \nu_- K : \|V\| = 1\},$$

the unit disk and unit sphere bundles in the negative normal bundle $\nu_- K$. then Theorem B has the following generalization:

Theorem B*. *If $[a, b]$ contains a single critical value c for F, the set of critical points with value c forming a nondegenerate critical submanifold K of Morse index λ, then $\mathcal{M}^b$ is homotopy equivalent to $\mathcal{M}^a$ with the disk bundle $D(\nu_- K)$ attached to $\mathcal{M}^a$ along $S(\nu_- K)$.*

The proof is a straightforward modification of the argument for Theorem B.

The above theorem gives an isomorphism on cohomology

$$H^k(\mathcal{M}^b, \mathcal{M}^a; \mathbb{Z}) \cong H^k(D(\nu_- K), S(\nu_- K); \mathbb{Z}).$$

When the bundle $\nu_- K$ is orientable and the group G preserves orientation of the normal bundle, we can use the Thom isomorphism theorem to evaluate the right-hand side:

$$H^k(D(\nu_- K), S(\nu_- K); \mathbb{Z}) \cong H^{k-\lambda}(K; \mathbb{Z}),$$

where λ is the Morse index of K.

We are interested in applying this theory to the case where $M = S^n$, the standard n-sphere, which we provide with the constant curvature metric, normalized so all sectional curvatures are one. Within M we have the prime totally geodesic two-spheres $M_{4\pi}$, a submanifold in which all elements have Morse index $n-2$. For α close to one, these totally geodesic submanifolds are also critical points for the α-energy E_α and $M_{4\pi}$ is a nondegenerate critical submanifold of Morse index $n-2$. The action of $O(3)$ on this submanifold is free and the quotient is diffeomorphic to the Grassmann manifold $G_3(\mathbb{R}^{n+1})$ of three-planes in $\mathbb{R}^{n+1}$. Moreover the negative normal bundle $\nu_- M_{4\pi}$ is trivial, hence orientable for any coefficient ring R.

Lemma 8.2. *There exists $\alpha > 1$ arbitrarily close to one such that any critical point for E_α either lies in $M_{4\pi}$ or has Morse index $\geq 2n-4$.*

This follows from the Nonbubbling Lemma 1.1, estimates of Ejiri [11] for prime minimal two-spheres, Proposition 7.2 for double covers of prime minimal two-spheres and Theorem 7.1 for branched covers of order $d \geq 3$.

Theorem 8.3. *We have isomorphisms*

$$H^k_{O(3)}(Map(S^2, S^n), S^n_0, R) \cong H^{k-n+2}_{O(3)}(M_{4\pi}; R)$$
$$\cong H^{k-n+2}(G_3(\mathbb{R}^{n+1}); R), \quad \text{for } k \leq 2n-5.$$

This follows from $O(3)$-equivariant Morse theory presented above because all α-energy critical points that are not in M_0 or $M_{4\pi}$ have Morse index $\geq 2n-4$.

Note in particular that the $H^*(BO(3); R)$-action on $H^k_{O(3)}(\mathrm{Map}(S^2, S^n), S^n_0, R)$ is trivial for $k \leq 2n-5$. We have calculated the low-dimensional equivariant cohomology using the constant curvature metric, but of course this is independent of the choice of Riemannian metric on S^n.

When S^n has the metric of constant curvature one, here are no minimal two-spheres $f : S^2 \to S^n$ of Morse index $\leq 2n-5$ which cover minimal projective planes $f_0 : \mathbb{R}P^2 \to S^n$. This follows from the results of Ejiri already cited; The minimal projective planes of lowest area and lowest Morse index in S^n are the constant curvature Veronese surfaces $f : S^2 \to S^4 \subseteq S^n$ which have Morse index 10 in S^4 and always have Morse index $\geq 2n-4$.

9 The Morse-Witten complex

Now that we know the equivariant cohomology of low degree, we want to use this to prove existence of minimal two-spheres on Morse index $\leq 2n-5$ in manifolds which satisfy our curvature hypothesis (7). The modern point of view is to organize this via a Morse-Witten complex, and we now review that idea.

Suppose first that M is a finite-dimensional smooth manifold, that $F : M \to [0, \infty)$ is a proper Morse function and that X is a "gradient-like" vector field for F, as defined in §3 of [24]. (For example, we might take X to be the gradient of F with respect to a Riemannian metric on M.) We choose an orientation for each critical point, that is, we choose an orientation of its unstable manifold, and let $C_m(F, -X)$ denote the free abelian group generated by the oriented critical points $\mathrm{Crit}_m(F)$ of F of Morse index m. If we choose X so that the stable and unstable manifolds of its critical points have transverse intersection (which is true for a residual set of X), we can define a boundary operator

$$\partial : C_m(F, -X) \to C_{m-1}(F, -X) \quad \text{by}$$
$$\partial(p) = \sum_{q \in \mathrm{Crit}_{m-1}(F)} n(p, q)q, \quad \text{for } p \in \mathrm{Crit}_m(F),$$

where $n(p, q)$ is the signed number of trajectories for $-X$ from x to y. Construction of this boundary is implicit in Milnor's proof of the h-cobordism theorem [24], and the theorems of standard Morse theory show that $\partial \circ \partial = 0$, and that the resulting chain complex

$$\longrightarrow C_{m+1}(F, -X) \longrightarrow C_m(F, -X) \longrightarrow C_{m-1}(F, -X) \longrightarrow \tag{41}$$

computes the homology of M. Of course, we can dualize and construct a corresponding cochain complex

$$\longrightarrow C^{m-1}(F, -X) \longrightarrow C^m(F, -X) \longrightarrow C^{m+1}(F, -X) \longrightarrow,$$
$$\text{where} \quad C^m(F, -X) = \mathrm{Hom}(C_m(F, -X), \mathbb{Z}). \tag{42}$$

We will call either $C_*(F, -X)$ or $C^*(F, -X)$ the *Morse-Witten complex* for F and X. More details of the resulting Morse homology can be found in the excellent treatment by Schwartz [33], although in the finite-dimensional case the analysis in his arguments can be replaced by the easier arguments of [24].

We can use the Morse theory on Banach manifolds of § 4 to extend the construction of the Morse-Witten complex to a perturbation of E_α which has nondegenerate critical points. In fact, as shown in [29], we can take the perturbation to be

$$E_{\alpha,\psi}(f) = E_\alpha(f) + \int_{S^2} f \cdot \psi dA, \tag{43}$$

where ψ is a suitable element of $L^2_k(S^2, \mathbb{R}^N)$. Condition C still holds for this perturbation within the completion $L^{2\alpha}_1(S^2, M)$, and implies that there are only finitely many critical points for $E_{\alpha,\psi}$ within

$$\mathcal{M}^{a-} = \{f \in L^2_k(S^2, M) : E_{\alpha,\psi}(f) < a\},$$

for any choice of $a \in \mathbb{R}$. We can then choose the gradient-like vector fields $\mathcal{X}$ so that stable and unstable manifolds intersect transversely, which enable us to define the Morse-Witten complex as in the finite-dimensional case [29].

But we can also consider the case of a perturbation of E_α which is equivariant Morse, all nonconstant critical points lying on isolated nondegenerate orbits for the symmetry group, in our case either $\hat{G} = O(3)$ or its identity component $G = SO(3)$. By Theorem 2.6 in [32] there is a small $\varepsilon > 0$ such that

$$\mathcal{M}^\varepsilon = L^2_k(S^2, M)^\varepsilon = \{f \in L^2_k(S^2, M) : E_\alpha(f) \leq \varepsilon\}$$

possesses the space M_0 of constant maps as a strong deformation retract.

Theorem 9.1. *The equivariant function E_α, restricted to $\mathcal{M}^{a-} - \mathcal{M}^\varepsilon$ for any $a > 0$ and a suitably small $\varepsilon > 0$, can be perturbed to an $SO(3)$-equivariant Morse function which is C^2 close to E_α and satisfies Condition C on the competion $L^{2\alpha}_1(S^2, M)$.*

Proof: This would follow from the Density Lemma 4.8 from Wasserman's treatment of equivariant Morse theory [38], except that our G-manifolds are not compact, and we want to ensure that our perturbed functions have smooth critical points and satisfy Condition C on the completion $L^{2\alpha}_1(S^2, M)$. We will explain how to modify Wasserman's argument so that it applies to our situation. We will assume known the theory of slices and tubes from Chapter 2 of [9], which carries over to infinite-dimensional G-manifolds with almost no change.

Note that by Lemma 3.1 we can assume that the isotropy group of E_α at each critical point is finite, and the isotropy is also finite on a small neighborhood of the compact critical locus for E_α within the open set $\mathcal{M}^{a-} - \mathcal{M}^\varepsilon$. If we make a C^2 small perturbation, the critical locus will remain in this neighborhood.

We give $\mathcal{M}^{a-} - \mathcal{M}^\varepsilon$ an increasing filtration by open sets

$$\mathcal{U}_n = \{f \in \mathcal{M}^{a-} - \mathcal{M}^\varepsilon : \text{ the isotropy group } G_f \text{ has order} \leq n \},$$

where $n \in \mathbb{N} \cup \{\infty\}$. We will prove by induction on n that for $n \in \mathbb{N}$ the restriction of E_α to $\mathcal{U}_n$ can be perturbed to a G-equivariant Morse function. We will do this by constructing a family of perturbed functions,

$$E_{\alpha,F}(f) = E_\alpha(f) + F(f), \quad \text{for} \quad F \in \mathcal{F}_n \subseteq C^2(L^2_k(S^2, M), \mathbb{R}),$$

to which we can apply the Baire category theorem. In fact the perturbations $\mathcal{F}_n$ will be a uniformly closed subspace of the space of C^2 functions

$$F : L^2_k(S^2, M) \longrightarrow \mathbb{R}$$

with the *strong topology* as described in §4 of Chapter 2 of [17]. As a basis for the open sets of this topology, we can take

$$\mathcal{N}(F, \varepsilon) = \{G \in C^2(L^2_k(S^2, M), \mathbb{R}) : d((j_2F)(f), (j_2G)(f)) < \varepsilon(f)\},$$

where j_2 denotes the two-jet and $\varepsilon : L^2_k(S^2, M) \to (0, 1)$ is an arbitrary positive continuous function. This is a Baire space by Theorem 4.2 of Chapter 2 of [17], so that the intersection of countably many open dense subsets of $\mathcal{F}_n$ is dense within $\mathcal{F}_n$.

Start of induction $n = 1$**:** The action of G on $\mathcal{U}_1$ is free and just as in the finite-dimensional case the quotient $\mathcal{U}_1/G$ is a smooth manifold, in fact, $\mathcal{U}_1$ is a principal G-bundle over $\mathcal{U}_1/G$. Since $L^2_k(S^2, M)$ has C^2 partitions of unity, we can construct countable open covers $\{U_{1,i} : i \in \mathbb{N}\}$ and $\{V_{1,i} : i \in \mathbb{N}\}$ of $\mathcal{U}_1$ consisting of G-invariant subsets with $\bar{V}_{1,i} \subseteq U_{1,i}$, together with a G-invariant partition of unity $\{\eta_{1,i} : i \in \mathbb{N}\}$ subordinate to $\{U_{1,i} : i \in \mathbb{N}\}$ with the support of $\eta_{1,i}$ contained within $\bar{V}_{1,i}$. Moreover, we can arrange that each $U_{1,i}$ contains a local slice $\mathcal{S}_{f_{1,i}}$ centered at $f_{1,i} \in U_{1,i}$ such that each G-orbit in $U_{1,i}$ intersects $\mathcal{S}_{f_{1,i}}$ exactly once. We can then define a family of perturbed functions

$$E_{\alpha,F}(f) = E_\alpha(f) + \sum_i \eta_{1,i} F_{1,i}(f),$$
$$\text{where} \quad F_{1,i}(f) = \int_{S^2} f \cdot \psi_{1,i} dA, \quad \text{for } f \in \mathcal{S}_{f_{1,i}}, \quad (44)$$

with $\psi_{1,i} \in L^2_k(S^2, \mathbb{R}^N)$ and $F_{1,i}$ is extended to be G-invariant. We let

$$\mathcal{F}_1 = \left\{ \sum_i \eta_{1,i} F_{1,i} \right\},$$

a family of functions on $L^2_k(S^2, M)$, where the open covers $\{U_{1,i}\}$ and $\{V_{1,i}\}$ are fixed but the partition of unity varies subject to the conditions

$$\text{supp}(\eta_{1,i}) \subseteq \bar{V}_{1,i}, \quad \|\eta_{1,i}\|_{C^2} \leq C_0, \quad (45)$$

where C_0 is a given bound. Then $\mathcal{F}_1$ is closed in the strong topology, and the usual argument shows that the restriction of $E_{\alpha,F}$ to $\mathcal{U}_1$ is an equivariant Morse function for a generic choice of perturbation $F \in \mathcal{F}_1$.

The case of general $n > 1$**:** We suppose that $\mathcal{F}_{n-1}$ has been constructed, and we let

$$\mathcal{N}_n = \{f \in \mathcal{U}_n : \text{ the isotropy group } G_f \text{ has order } n \},$$

a smooth submanifold with $\mathcal{U}_n = \mathcal{U}_{n-1} \cup \mathcal{N}_n$.

We construct countable open covers $\{U_{n,i} : i \in \mathbb{N}\}$ and $\{V_{n,i} : i \in \mathbb{N}\}$ of $\mathcal{N}_n$, consisting of G-invariant open subsets of $\mathcal{U}_n$ with $\bar{V}_{n,i} \subseteq U_{n,i}$, together with a G-invariant partition of unity $\{\eta_{n,i} : i \in \mathbb{N}\}$ subordinate to $\{U_{n,i} : i \in \mathbb{N}\}$ with $\text{supp}(\eta_{n,i})$ contained within $\bar{V}_{n,i}$. We assume that $U_{n,i}$ contains $f_{n,i} \in \mathcal{N}_n$, and that $U_{n,i}$ possesses a local slice $\mathcal{S}_{f_{n,i}}$ centered at $f_{n,i}$, with each G-orbit within $U_{n,i}$ intersecting $\mathcal{S}_{f_{n,i}}$. We can suppose that $\mathcal{S}_{f_{n,i}}$ is the domain of a tangent-space-valued chart centered at $f_{n,i}$,

$$\phi_{n,i} : \mathcal{S}_{f_{n,i}} \longrightarrow E \oplus F = T_{f_{n,i}} \mathcal{S}_{f_{n,i}},$$
$$\text{where} \quad \phi_{n,i}(\mathcal{N}_n \cap \mathcal{S}_{f_{n,i}}) = (E \oplus \{0\}) \cap \phi_{n,i}(\mathcal{S}_{f_{n,i}}).$$

Indeed, according to Bochner's Theorem (see 2.2.1 in [9]) we can assume that the differential of $\phi_{n,i}$ is the identity, and that $\phi_{n,i}$ takes the action of the

isotropy group $G_{f_{n,i}}$ on $f_{n,i}$ to an orthogonal linear action on the tangent space which preserves the direct sum decomposition $E \oplus F$ and is trivial on E. We can assume also that $\phi_{n,i}(\mathcal{S}_{f_{n,i}})$ is a product neighborhood in $E \oplus F$.

We can think of $\phi_{n,i}$ as defining an E-valued chart on $\mathcal{N}_n \cap \mathcal{S}_{f_{n,i}}$, and as in the case $n = 1$ we can define a family of perturbed functions on $\mathcal{N}_n$,

$$E_{\alpha,F}(f) = E_\alpha(f) + \sum_i \eta_{n,i} F_{n,i}(f),$$
$$\text{where} \quad F_{n,i}(f) = \int_{S^2} \phi_{n,i}(f) \cdot \psi_{n,i} dA, \quad \text{for } f \in \mathcal{S}_{f_{n,i}}, \quad (46)$$

with $\psi_{n,i} \in E$, where E is a linear subspace of $L^2_k(S^2, \mathbb{R}^N)$ and $F_{n,i}$ is extended to be G-invariant on $U_{n,i}$. As in the case $n = 1$, the restriction to $U_{n,i} \cap \mathcal{N}_n$ will be Morse for a generic choice of $\psi_{n,i}$'s.

Finally, we let P_E and P_F denote orthogonal projections to E and F respectively, and define

$$r_i^2 : U_{n,i} \to [0, \infty) \quad \text{by} \quad r_i^2(f) = \|P_F(\phi_{n,i}(f))\|^2.$$

We then modify (46) to the family of perturbations

$$E_{\alpha,F}(f) = E_\alpha(f) + \sum_i \eta_{n,i} \left[\varepsilon_i r_i^2(f) + F_{n,i}(f)\right],$$
$$\text{where} \quad F_{n,i}(f) = \int_{S^2} P_E(\phi_{n,i}(f)) \cdot \psi_{n,i} dA, \quad \text{for } f \in \mathcal{S}_{f_{n,i}}, \quad (47)$$

Since r_i^2 is invariant under the action of the isotropy group $G_{f_{n,i}}$, we can extend $E_{\alpha,F}$ to $U_{n,i}$ so that it is G-invariant. For a generic choice of $\psi_{n,i}$'s, this will have the property that all critical points within $U_{n,i} \cap \mathcal{N}_n$ are nondegenerate.

To complete the inductive step we now set

$$\mathcal{F}_n = \mathcal{F}_{n-1} \cup \left\{ \sum_i \eta_{n,i} \left[\varepsilon_i r_i^2(f) + F_{n,i}(f)\right] \right\},$$

where the open covers $\{U_{n,i}\}$ and $\{V_{n,i}\}$ are fixed but the partition of unity varies subject to the conditions

$$\text{supp}(\eta_{n,i}) \subseteq \bar{V}_{n,i}, \quad \|\eta_{n,i}\|_{C^2} \leq C_0, \quad (48)$$

where C_0 is a given bound. Once again, the Baire category argument implies that the resulting perturbed functions will be Morse for generic elements of $\mathcal{F}_n$.

Finally, we observe that our perturbations do not involve the derivatives of the map f and can be made arbitrarily small. The fact that (48) holds for all $n \in \mathbb{N}$ makes it easy to check that Condition C still holds on the completion, and that the regularity argument of Sacks and Uhlenbeck implies that any critical point must be in $L^2_k(S^2, M)$. These results are discussed further in [29]. This finishes the proof.

Note that if M is C^∞, we can take k to be arbitrary in Theorem 9.1, and thus we can arrange that all critical points are C^∞.

We want to apply Theorem 9.1 to construct an equivariant Morse-Witten complex. Under the hypotheses of this theorem, all nonconstant minimal two-spheres of Morse index $\leq 2n-5$ in M are immersions and lie on Morse nondegenerate critical submanifolds, each a $\widehat{\mathcal{G}}$-orbit, where $\widehat{\mathcal{G}}$ is the group of conformal and anticonformal automorphisms of S^2 There are only finitely many of these by Theorem 6.3 and according to Theorem 2.2, these perturb to finitely many G-orbits of α-energy crticial points, each a nondegenerate critical submanifold. Let U be an open subset which contains all critical points of Morse index $\leq 2n-5$, and suppose that V is a larger open set such that $U \subseteq \bar{U} \subseteq V$ and V contains no additional critical points. Since $L_1^{2\alpha}(S^2, M)$ has C^2 partitions of unity, we can construct a smooth function $\eta : L_1^{2\alpha}(S^2, M) \to \mathbb{R}$ which is zero on U and one on $L_1^{2\alpha}(S^2, M) - v$. Finally, we let

$$F : L_1^{2\alpha}(S^2, M) \longrightarrow \mathbb{R} \quad \text{by} \quad F = (1-\eta)E_\alpha + \eta\tilde{F},$$

where $\tilde{F}$ is an equivariant Morse approximation to E_α as described in the previous paragraph, all of its critical points lying on Morse nondegenerate critical submanifolds, each an orbit for G.

Under the hypotheses of Theorems 1.1 and 1.3, all prime minimal two-spheres of Morse index $\leq 2n-5$ in M are immersions and lie on Morse nondegenerate critical submanifolds, each a $\widehat{\mathcal{G}}$-orbit, where $\widehat{\mathcal{G}}$ is the group of conformal and anticonformal automorphisms of S^2 There are only finitely many of these by Theorem 6.3 and according to Theorem 2.2, these perturb to finitely many G-orbits of α-energy criticial points, each a nondegenerate critical submanifold. Let U be an open subset which contains all prime critical points of Morse index $\leq 2n-5$, and suppose that V is a larger open set such that $U \subseteq \bar{U} \subseteq V$ and V contains no additional critical points. Since $L_1^{2\alpha}(S^2, M)$ has C^2 partitions of unity, we can construct a smooth function $\eta : L_1^{2\alpha}(S^2, M) \to \mathbb{R}$ which is zero on U and one on $L_1^{2\alpha}(S^2, M) - V$. Finally, we let

$$F : L_1^{2\alpha}(S^2, M) \longrightarrow \mathbb{R} \quad \text{by} \quad F = (1-\eta)E_\alpha + \eta\tilde{E}_\alpha,$$

where $\tilde{E}_\alpha$ is a nearby equivariant Morse function. Then F is equivariant Morse function which agrees with E_α on U.

Note that except for points lying within the space M_0 of constant maps, the isotropy groups of E_α are always finite by Lemma 3.1, and we can arrange that F satisfies the same condition. Let $\{\phi_t : t \in [0,\infty)\}$ be the one-parameter semigroup of diffeomorphisms of $\mathcal{M}$ corresponding to $-\mathcal{X}$, this semigroup being defined for all positive t because $-\mathcal{X}$ is positively complete. Note that the isotropy group for either the action of G or G_0 is constant along any orbit for $-\mathcal{X}$. Recall that the *unstable manifold* $W_p(F, \mathcal{X})$ of a critical point $p \in \mathcal{M}$ consists of the images of all trajectories $t \mapsto \phi_t(q)$ which start at f in the remote past, in other words, such that $\phi_t(q) \to p$ as $t \to -\infty$. Similarly, the *stable manifold* $W_p^*(F, \mathcal{X})$ of a critical point p consists of the images of trajectories $t \mapsto \phi_t(q)$ such that $\phi_t(q) \to p$ as $t \to \infty$.

Lemma 9.2. *The unstable and stable manifolds $W_p(f, \mathcal{X})$ and $W_p^*(f, \mathcal{X})$ are indeed submanifolds of $\mathcal{M}$.*

The proof follows from the explicit form of the gradient-like vector field $\mathcal{X}$ near f and the properties of local flows for vector fields on Banach manifolds.

It follows from Condition C and the explicit form of $\mathcal{X}$ that any orbit $g \mapsto \phi_t(g)$ converges to some critical point as $t \to \infty$ (although it may come close to several other critical points first).

Lemma 9.3. *We can adjust the G_0-equivariant gradient-like vector field $\mathcal{X}$ so that if λ_p is the Morse index of the critical point $p \in \mathcal{M}$ and λ_q is the Morse index of q,*

1. *$\lambda_p \leq \lambda_q \Rightarrow W_p(\mathcal{X}) \cap W_q^*(\mathcal{X})$ is empty, while*
2. *$\lambda_p > \lambda_q \Rightarrow W_p(\mathcal{X}) \cap W_q^*(\mathcal{X})$ is a submanifold of dimension $\lambda_p - \lambda_q + 3$.*

Proof: If $W_p(\mathcal{X}) \cap W_q^*(\mathcal{X})$ is nonempty, then $F(p) > F(q)$ and we can choose a regular value c for F such that $F(p) > c > F(q)$, noting that we can choose c as close to $F(p)$ as we want. Let $\mathcal{S}_p$ be a section for the G_0-action through the point p, a submanifold of codimension three.

Then $\mathcal{N} = F^{-1}(c)$ is a codimension one submanifold of $\mathcal{M}$, and

$$\text{dim } (\mathcal{N} \cap \mathcal{S}_p \cap W_p(\mathcal{X})) = \lambda_p - 1, \qquad \text{codim } (\mathcal{N} \cap W_q^*(\mathcal{X})) = \lambda_q.$$

If we choose c sufficiently close to the Morse nondegenerate critical point p, we see that $S = \mathcal{N} \cap \mathcal{S}_p \cap W_p(\mathcal{X})$ is a $(\lambda_p - 1)$-dimensional sphere and lies in an open tubular neighborhood U of $\mathcal{N}$ with diffeomorphism $\psi : U \to S \times V$, where V is an open ball in a Banach space, with $\psi(S) = S \times \{0\}$. Let

$$N = U \cap W_q^*(\mathcal{X}), \quad \text{and consider} \quad G = \pi \circ \psi : N \to V,$$

where π is the projection on the second factor. We can check that g is a Fredholm map with Fredholm index $\lambda_p - 1 - \lambda_q$, and

$$\psi^{-1}(S \times \{x\}) \cap N = \{s \in N : g(s) = x\}.$$

Assuming that $\mathcal{X}$ (and hence N) is sufficiently smooth, we can use the Sard-Smale Theorem of [36] to choose a regular value x for g. Finally, we can choose x as close as we want to 0, and construct an isotopy from a neighborhood of S to itself which carries $S \times \{0\}$ to $S \times \{x\}$. Since the conditions defining pseudogradient are open, we can replace $\mathcal{X}$ by a new gradient-like vector field so that

$$\mathcal{N} \cap \mathcal{S}_p \cap W_p(\mathcal{X}) \quad \text{and} \quad \mathcal{N} \cap W_q^*(\mathcal{X})$$

have transverse intersection. Then the intersection of

$$\mathcal{N} \cap W_p(\mathcal{X}) \quad \text{and} \quad \mathcal{N} \cap W_q^*(\mathcal{X})$$

will be a submanifold of dimension $\lambda_p - \lambda_q + 3$.

Since F satisfies Condition C, there are only finitely many G-orbits within

$$\mathcal{M}^a = \{f \in L_1^{2\alpha}(S^2, M) : F(f) \leq a\},$$

for any choice of $a \in \mathbb{R}$. Lemma 9.2 implies that after perturbation of $-\mathcal{X}$ unstable manifolds of critical orbits can only intersect stable manifolds of critical points of lower index. Using the arguments in [24], we can then modify the function F to a new function $\tilde{F}$, without changing G-orbits of critical points or trajectories between critical points, so that $\tilde{F}$ is a self-indexing equivariant Morse function, one for which each critical point $f \in \mathcal{M}^a$ satisfies

$$\tilde{F}(f) = \text{Morse index of } f.$$

Now we can use further arguments within [24] to define the Morse-Witten complexes (41) and (42). Let us focus on the cohomology complex. For $\lambda \in \{0\} \cup \mathbb{N}$, we let

$$C^\lambda(F, -\mathcal{X}) = H_G^\lambda(\mathcal{M}^{\lambda+(1/2)}, \mathcal{M}^{\lambda-(1/2)}; \mathbb{Z}),$$

and define

$$d : C^\lambda(F, -\mathcal{X}) \longrightarrow C^{\lambda+1}(F, -\mathcal{X})$$

to be the coboundary operator in the triple

$$(\mathcal{M}^{\lambda+(3/2)}, \mathcal{M}^{\lambda+(1/2)}, \mathcal{M}^{\lambda-(1/2)}).$$

using the fact that the cohomology of the pair $(\mathcal{M}^{\lambda+(1/2)}, \mathcal{M}^{\lambda-(1/2)})$ is concentrated in degree λ, we then obtain a cochain complex

$$\longrightarrow C^{\lambda-1}(\tilde{F}, -\mathcal{X}) \longrightarrow C^\lambda(\tilde{F}, -\mathcal{X}) \longrightarrow C^{\lambda+1}(\tilde{F}, -\mathcal{X}) \longrightarrow,$$

which we call the Morse-Witten complex, in which each of the coboundary maps d is an $H^*(BG)$-module homomorphism. In the case where the action of G is free, this does indeed count trajectories connecting critical points in the orbit space $\mathcal{M}/G$, and in keeping with the philosophy of equivariant cohomology, we regard this as counting trajectories in $\mathcal{M}/G$ even when the action is only locally free.

10 Existence of low index minimal two-spheres

Finally, we collect the preceding ingredients to prove existence of minimal two-spheres of Morse index $\leq 2n - 5$ in manifolds which satisfy the curvature conditions (2) or (7), and thus prove Theorems 1.1 and 1.3.

Suppose that M is homeomorphic to S^n and that M is given a Riemannian metric of positive half-isotropic sectional curvatures. Then it follows from Theorem 6.2 that there are only finitely many $\widehat{\mathcal{G}}$-orbits of prime minimal two spheres

$f : S^2 \to M$ with Morse index $\leq 2n-5$, where $\widehat{\mathcal{G}} = PSL(2,\mathbb{C}) \cup R \cdot PSL(2,\mathbb{C})$. Some of these may be invariant under an antipodal map A, and hence double cover minimal projective planes. If $\alpha_0 > 1$ is sufficiently close to one, each of these $\widehat{\mathcal{G}}$-orbits perturbs to a single nondegenerate $O(3)$-orbit of critical points for E_α for $\alpha \in (1, \alpha_0)$. Moreover, for Morse index $\leq 2n-5$, there are no sequences of α_m-energy critical points with $\alpha_m \to 1$ which approach bubble trees with more than one bubble by Theorem 1.1. Indeed, if we have a sequence of such α_m-energy critical points with $\alpha_m \to 1$, a subsequence must either converge to an orbit of prime minimal two-spheres or a branched cover of such a prime minimal two spheres. Under the curvature condition (7), any such branched cover must be of order two, since otherwise it would have index $\geq 2n-4$ by Theorem 7.1.

After imposing the center of mass condition described in §2, parametrized minimal two-spheres $f : S^2 \to M$ of Morse index $< 2n-4$ are partitioned into several types depending upon their isotropy groups for the action of $\widehat{G} = O(3)$:

1. We have prime minimal two-spheres for which the isotropy group of the $\widehat{G}$-action is trivial. In this case, the $\widehat{G}$-orbit consists of two components, one with each of two orientations.

2. We may also have prime minimal projective planes for which the isotropy group H of the $\widehat{G}$-action is isomorphic to $\mathbb{Z}_2$ and is generated by an orientation-reversing element $A \in \widehat{G}$. In this case, the $\widehat{G}$-orbit consists of a single component.

3. Next we may have nontrival branched covers of prime minimal two-spheres, for which the isotropy group H of $\widehat{G}$ is a nontrivial discrete subgroup of $G = SO(3)$. Under our curvature hypothesis, the branching order must be two.

4. Finally, we might have nontrivial branched covers of prime minimal projective planes.

Since bubbling is ruled out, sequences of α_m-energy critical points with $\alpha_m \to 1$ must converge to a minimal two-sphere which lies in one of these cases. We claim that the cases can be distinguished within equivariant cohomology.

We now construct a Morse-Witten complex for the function E_α on the pair $(\mathcal{M}, M_0)$ as in § 9, and determine the contributions given by Theorem B* of §8 to the cohomology of this complex when all critical $\widehat{G}$-orbits are nondegenerate. We claim that the type of critical orbit obtained is encoded in the action of

$$H^*(BO(3); \mathbb{Z}_2) = P[w_1, w_2, w_3]$$

on the equivariant cohomology of the corresponding generator in the Morse-Witten complex. Suppose we want to calculate the contribution to the Morse-Witten complex of an isolated nondegenerate critical orbit K with isotropy group H, so K is diffeomorphic to $\widehat{G}/H$. The standard approach for computing

the equivariant cohomology of K is to use the spectral sequence of the fibration

$$\begin{array}{ccc} K & \longrightarrow & K_{\widehat{G}} \\ & & \pi\downarrow \\ & & B\widehat{G}, \end{array} \tag{49}$$

the equivariant cohomology of K being the usual cohomology of the total space $K_{\widehat{G}}$.

1. In the case of trivial isotropy, $K = O(3)$ and the homotopy quotient is contractible, and this forces the action of $P[w_1, w_2, w_3]$ to be trivial.

2. In the case of prime minimal projective planes, in which the isotropy group is isomorphic to $\mathbb{Z}_2$ and is generated by an orientation-reversing element $A \in O(3)$, the orbit space is $K = SO(3)$ which has $\mathbb{Z}_2$ cohomology

$$H^*(SO(3); \mathbb{Z}_2) \cong \Lambda(\xi_1, \xi_2), \quad \text{where} \quad \deg \xi_i = i,$$

and the spectral sequence shows that w_1 acts freely on the homotopy quotient $K_{\widehat{G}}$, while w_2 and w_3 act trivially.

3 and 4. For the remaining cases, we consider the isotropy group under the action of $G = SO(3)$. For double covers, the isotropy group is $\mathbb{Z}_2$ and is generated by a rotation, while the orbit space $K = SO(3)/\mathbb{Z}_2$ has fundamental group $\pi_1(K) \cong \mathbb{Z}_4$. It follows from Poincaré duality that

$$H^*(K; \mathbb{Z}_2) \cong \Lambda(\eta_1, \eta_2), \quad \text{where} \quad \deg \xi_i = i,$$

once again, but the projection $\pi : SO(3) \to K$ induces a map on $\mathbb{Z}_2$-cohomology which takes η_1 to zero. This implies that in the spectral sequence of (49), w_2 is not in the image of transgression, which implies that w_2 must act freely on the homotopy quotient in this case. Since the equivariant cohomology of K is just $H_{SO(3)}(K; \mathbb{Z}_2) \cong P[u]$, a polynomial algebra generated by an element u of degree one, it must be that w_2 acts by taking 1 and u to $1 \cdot w_2 = u^2$ and $uw_2 = u^3$ respectively, and in particular the action of $O(3)$ is nontrivial.

Now we use Theorem B* of §8 and the Thom isomorphism theorem to show that each critical orbit yields a generator for

$$H^*_{O(3)}(\mathcal{M}, M_0, \mathbb{Z}_2) \quad \text{as a module over} \quad P[w_1, w_2, w_3]$$

the degree of the generators being shifted upward by the Morse index of the critical orbit.

We now divide the Morse-Witten complex $C^\lambda(F, -\mathcal{X})$ into a direct sum,

$$C^\lambda(F, -\mathcal{X}) = A^\lambda(F, -\mathcal{X}) \oplus B^\lambda(F, -\mathcal{X}),$$

where $H^*(BO(3); \mathbb{Z}_2) = P[w_1, w_2, w_3]$ acts trivially on A^λ and nontrivially on B^λ, the generators of A^λ being prime minimal two-spheres with two possible orientations, the generators of B^λ being projective planes and double covers of prime minimal two-spheres or projective planes. Note that

$$P[w_1, w_2, w_3] \text{ acts trivially on } A^\lambda \quad \Rightarrow \quad d(A^\lambda) \subseteq A^\lambda,$$

because the differential d commutes with the action of $P[w_1, w_2, w_3]$. Thus the A^λ's form a subcomplex of the Morse-Witten complex, and we have a short exact sequence

$$0 \longrightarrow A^* \longrightarrow C^* \longrightarrow B^* \longrightarrow 0.$$

But we have noted before that in the constant curvature case, there are no double covers of real projective planes with Morse index $\leq 2n-5$, so we have

$$H^k(A^*; \mathbb{Z}_2) \cong H^k(C^*; \mathbb{Z}_2) \cong H^{k-n+2}(G_3(\mathbb{R}^{n+1}); \mathbb{Z}_2),$$

when $k \leq 2n-5$, the last isomorphism following from Theorem 8.3. Note that the $\widehat{\widehat{G}}$ action does not depend on the Riemannian metric so we must have the same result for metrics satisfying our curvature hypotheses. The Morse inequalites for the subcomplex A^* now shows that the number of minimal two-spheres which do not cover projective planes of Morse index λ for $n-2 \leq \lambda \leq 2n-5$ must be at least as large as the number $p_3(\lambda - n + 2)$, where $p_3(k)$ is the number of k-cells in the Schubert cell decomposition for $G_3(\mathbb{R}^{n+1})$, which proves the Main Theorem 1.1, and its extension to the weaker curvature hypothesis (7).

References

[1] D. M. Austin and P. J. Braam, *Morse-Bott theory and equivariant cohomology*,

[2] W. Ballmann, G. Thorbergsson and W. Ziller, *Existence of closed geodesics on positively curved manifolds*, J. Diff. Geometry **18** (1983), 221-252.

[3] M. Berger, *Sur quelques variétés riemanniennes suffisamment pincées*, Bull. S. M. F. **88** (1960), 57-71.

[4] R. Bott, *Lectures on Morse theory, old and new*, Bull. Amer. Math. Soc. **7** (1982), 331-358.

[5] E. Calabi, *Minimal immersions of surfaces in Euclidean spheres*, J. Differential Geometry **1** (1967), 111-125.

[6] J. Chen and G. Tian, *Compactification of moduli space of harmonic mapping*, Comm. Math. Helv. **74** (1999), 201-237.

[7] S. S. Chern, *on minimal spheres in the four sphere*, Studies and essays presented to Y. W. Chen, Taiwan (1970), 137-150.

[8] H. Choi and R. Schoen, *The space of minimal embeddings of a surface into a three-dimensional manifold of positive Ricci curvature*, Inventiones math. **81** (1975), 387-394.

[9] J. Duistermaat and J. Kolk, *Lie groups*, Springer, New York, 2000.

[10] D. Ebin, *The manifold of Riemannian metrics*, Proc. Symposia in Pure Math., Amer. Math. Soc. **15**, 11-40.

[11] N. Ejiri, *The index of minimal immersions of S^2 into S^{2n}*, Math. Z. **184** (1983), 127-132.

[12] D. Gromoll and W. Meyer, *On differentiable functions with isolated critical points*, Topology **8** (1969), 361-369.

[13] M. Gromov, *Homotopical effects of dilatation*, J. Differential Geometry **13** (1978), 303-310.

[14] A. Grothendieck, *Sur la classification des fibres holomorphes sur la sphere de Riemann*, Amer. J. Math. **79** (1957), 121-138.

[15] R. Gunning, *Lectures on Riemann surfaces*, Princeton Univ. Press, Princeton, NJ, 1966.

[16] N. Hingston, *Equivariant Morse theory and closed geodesics*, J. Differential Geometry **19** (1978), 85-116.

[17] M. Hirsch, *Differenital topology*, Springer, New York, 1976.

[18] N. Korevaar, *Lower bounds for eigenvalues of conformal metrics*, J. Differential Geometry **37** (1993), 73-93.

[19] S. Lang, *Differential and Riemannian manifolds*, Springer, New York, 1996.

[20] B. Lawson and M. Michelsohn, *Spin geometry*, Princeton Univ. Press Princeton, NJ, 1989.

[21] P. Li and S. T. Yau, *On the Schrödinger equation and the eigenvalue problem*, Comm. Math. Physics **88** (1983), 309-318.

[22] M. Micallef and J. D. Moore, *Minimal two-spheres and the topology of manifolds with positive curvature on totally isotropic two-planes*, Annals of Math. **127** (1988), 199-227.

[23] J. Milnor, *Morse theory*, Annals of Math. Studies **51**, Princeton Univ. Press, Princeton, NJ, 1963.

[24] J. Milnor, *Lectures on the h-cobordism theorem*, Princeton Univ. Press, Princeton, NJ, 1965.

[25] J. Milnor, *Geometry and dynamics of quadratic rational maps*, Experiment. Math. **2** (1993), 37-83.

[26] J. Milnor and J. Stasheff, *Characteristic classes*, Princeton Univ. Press, Princeton NJ, 1974.

[27] S. Montiel and F. Urbano, *Second variation of superminimal surfaces into self-dual Einstein four-manifolds*, Trans. AMS **349** (1997), 2253-2267.

[28] J. D. Moore, *On the number of minimal two-spheres of small area in manifolds with curvature bounded above*, Math. Z. **288** (1990), 323-343.

[29] J. D. Moore, *Lectures on global analysis: minimal surfaces in Riemannian manifolds*, to appear.

[30] R. Palais, *Lusternik-Schnirelmann theory on Banach manifolds*, Topology **5** (1966), 115-132.

[31] T. Parker, *Bubble tree convergence for harmonic maps curvature flow*, J. Differential Geometry **44** (1996), 595-633.

[32] J. Sacks and K. Uhlenbeck, *The existence of minimal immersions of* 2*-spheres*, Annals of Math. **113** (1981), 1-24.

[33] M. Schwarz, *Morse homology*, Birkhäuser Verlag, Basel, 1993.

[34] G. Segal, *The topology of spaces of rational functions*, Acta Mathematica **143** (1979), 39-72.

[35] S. Smale, *Morse theory and a nonlinear generalization of the Dirichlet problem*, Annals of Math. **80** (1964), 382-396.

[36] S. Smale, *An infinite-dimensional version of Sard's theorem*, Amer. J. Math. **87** (1966), 861-866.

[37] K. Uhlenbeck, *Morse theory on Banach manifolds*, J. Functional Analysis **10** (1972), 430-445.

[38] A. G. Wasserman, *Equivariant differential topology*, Topology **8** (1969), 127-150.

[39] B. White, *Every three-sphere of positive Ricci curvature contains a minimal imbedded torus*, Bull. Amer. Math. Soc. **21** (1989), 71-75.

[40] B. White, *The space of minimal submanifolds for varying Riemannian metrics*, Indiana Math. J. **40**, (1991), 161-200.

[41] J. Wolf, *Spaces of constant curvature*, McGraw-Hill, New York, 1967.

[42] P. Yang and S. T. Yau, *Eigenvalues of the Laplacian of compact Riemann surfaces and minimal submanifolds*, Annali Scuola Normale Superiore Pisa, **7** (1980), 55-63.